

2023

SHARP Survey

(Prevention Needs Assessment)
Results for White Students

State of Utah

Department of Health & Human Services
Office of Substance Use and Mental Health



TABLE OF CONTENTS

Acknowledgments 3
Introduction 4
Understanding the Charts 5
Substance Use 9
Substance Use 10
Antisocial Behavior 18
Community and School Climate and Safety 19
Social and Emotional Health 24
Physical Health and Safety 38
The Risk and Protective Factor Model of Prevention 42
Risk and Protective Factors 44
School and Community Improvement Using Survey Data 56
Building a Strategic Prevention Framework 57
Appendix A: Changes between PNA administrations 59
Appendix B: Substance Use and Perceived Parental Acceptability 61
Appendix C: Drug Free Communities Data 62
Appendix D: Contacts for Prevention 63
Appendix E: Risk and Protective Scale Definitions 67
Appendix F: Item Dictionary 70

ACKNOWLEDGMENTS

Coordination and administration of the Utah SHARP/PNA Survey was a collaborative effort of the State of Utah, Department of Health and Human Services; and Bach Harrison, LLC. For more information about the SHARP/PNA or prevention services in Utah, please refer to Appendix D: Contacts for Prevention.

INTRODUCTION

2023 White Students Prevention Needs Assessment Survey Report

In Spring 2023, the Utah Department of Health and Human Services (DHHS) conducted the School Health and Risk Prevention (SHARP) survey project in public schools throughout Utah. This project included the Prevention Needs Assessment (PNA), the results of which are presented in this report. The PNA surveyed students in grades 6, 8, 10, and 12 on substance abuse, mental health, chronic conditions, healthy lifestyles, violence and injury, social determinants of health, and risk and protective factors. The survey was administered by a monitored web survey or a self-administered paper/pencil questionnaire. Across the State,

35 school districts, 13 charter schools, and 1 private school participated in SHARP/PNA.

Results across seven categories are presented in this SHARP/PNA report: substance use, antisocial behavior, community and school climate and safety, social and emotional health, substance-related indicators, physical health and safety, and risk and protective factors.

Please see the Characteristics of Participants table below for participation information, and note that because not all students answer all of the questions, the total number of survey respondents by gender and survey respondents by ethnicity may be less than the reported total students.

Characteristics of participants

	Race	2010 l	Race	2021	Race	2023	State 2023				
	Number	Percent	Number	Percent	Number	Percent	Number	Percent			
What grade are you in	n?										
6	19,493	31.2	17,085	32.1	11,617	31.6	17,512	33.7			
8	18,400	29.4	15,981	30.1	11,757	32.0	16,525	31.8			
10	15,123	24.2	12,746	24.0	8,933	24.3	11,997	23.1			
12	9,510	15.2	7,340	13.8	4,445	12.1	5,856	11.3			
All grades	62,526	100.0	53,152	100.0	36,752	100.0	51,890	100.0			
Are you:											
Male	29,645	47.6	25,709	48.7	17,830	49.0	25,042	48.8			
Female	32,135	51.6	27,070	51.3	18,572	51.0	26,299	51.2			
Which of the followin	g best desc	cribe you?*	•								
Heterosexual (straight)	37,675	89.1	29,675	84.5	18,620	82.7	25,157	81.3			
Gay or lesbian	585	1.4	655	1.9	551	2.4	811	2.6			
Bisexual	1,987	4.7	2,471	7.0	1,700	7.6	2,526	8.2			
Not sure/Other	2,056	4.9	2,311	6.6	1,632	7.3	2,449	7.9			
What is your race? (Se	elect one o	r more)†									
African American	0	0.0	0	0.0	0	0.0	775	1.5			
American Indian	0	0.0	0	0.0	0	0.0	619	1.2			
Asian	0	0.0	0	0.0	0	0.0	831	1.6			
Hispanic or Latino	0	0.0	0	0.0	0	0.0	8,290	16.3			
Pacific Islander	0	0.0	0	0.0	0	0.0	616	1.2			
White	62,526	100.0	53,152	100.0	36,752	100.0	36,752	72.4			
Multi-racial	0	0.0	0	0.0	0	0.0	2,869	5.7			

^{*} A question about sexual orientation was added to the 2019 survey instrument used by 8th, 10th, and 12th graders. (6th grade surveys omitted this question.) If any response category has fewer than 10 respondents, all question data are suppressed to protect student privacy.

[†] To accurately represent Hispanic/Latino participation in the SHARP survey, students indicating they are of Hispanic or Latino ethnicity and up to one race are reported as Hispanic or Latino. Students reporting more than one race are reported as multi-racial (regardless of ethnicity).

^{10/10/2023:} We have removed 44 surveys from students on the Navajo Nation that were included in the first version of this report. This data is property of the Navajo Nation and we apologize for not catching the error sooner. This has an incredibly minimal impact on the data trends for the state and regions.

UNDERSTANDING THE CHARTS

There are seven types of charts presented in this report:

- 1. Substance use
- 2. Antisocial behavior
- 3. Community and school climate and safety
- 4. Social and emotional health
- 5. Substance-related indicators
- 6. Physical health and safety
- 7. Risk factor profiles
- 8. Protective factor profiles

Data from the charts are presented in tables directly below them. Additional data useful for prevention planning are found in Appendices B and C. Note that data reported in the tables are rounded to one decimal place. (Rates of 0% to 0.049% are displayed as 0.0%.)

Understanding the Format of the Charts

There are several graphical elements common to all the charts. Understanding the format of the charts and what these elements represent is essential in interpreting the results of the 2023 SHARP survey.

The bars found in most of the charts, for example, on substance use and antisocial behavior charts, represent the percentage of students in that grade who reported a given behavior. The bars on the risk and protective factor charts represent the percentage of students whose answers reflect significant risk or protection in that category.

Each set of differently colored bars represents one of the last three administrations of the SHARP (PNA): 2019, 2021, and 2023. By looking at the percentages over time, it is possible to identify trends in substance use and antisocial behavior. By studying the percentage of youth at risk and with protection over time, it is possible to determine whether the percentage of students at risk or with protection is increasing, decreasing, or staying the same. This information is important when deciding which risk and protective factors warrant attention.

Dots, Diamonds, Triangles, and Xs provide points of comparison to larger samples. The dots on the charts represent the

percentage of all of the youth surveyed across Utah who reported substance use, problem behavior, elevated risk, or elevated protection. The diamonds and stars represent national data from the Monitoring the Future (MTF) Survey and the Bach Harrison Norm, respectively.

The dots represent state-level data for the current year. For the 2023 SHARP (PNA) Survey, there were 51,934 participants in grades 6, 8, 10, and 12. Data were cleaned and weighted by school, grade, and gender to ensure that the data reported at all levels accurately represent the population of Utah. The survey results provide considerable information for communities to use in planning prevention services.

A comparison to state-wide and national results provides additional information for your community in determining the relative importance of levels of Alcohol, Tobacco, and Other Drugs (ATOD) use, antisocial behavior, risk, and protection. Information about other students in the state and the nation can be helpful in determining the seriousness of a given level of problem behavior.

Note that while the national data are visually represented on the charts, they may be omitted from the tables due to limitations on available space in the new SHARP (PNA) report format.

The diamonds represent national data from the Monitoring the Future (MTF) survey, a long-term epidemiological study that surveys trends in drug and alcohol use among American adolescents. Funded by research grants from the National Institute on Drug Abuse, it features nationally representative samples of 8th-, 10th-, and 12th-grade students. (MTF is not administered to 6th graders and as such 6th grade data cannot be represented on the charts.)

The triangles represent national data from the Bach Harrison Norm (BH Norm). Bach Harrison Norm was developed by Bach Harrison LLC to provide states and communities with the ability to compare their results on risk, protection, and antisocial measures with more national measures. Survey participants from 11 statewide surveys were combined into a database of approximately 657,000 students in grades 6, 8, 10, and 12. The results were weighted to make the contribution of each state proportional to its share of the national population. Bach Harrison analysts then calculated rates for antisocial behavior and for students at risk and with protection. The results appear on the charts as the BH Norm. In order to keep the Bach Harrison Norm

relevant, it is updated approximately every 2 years as new data become available. An update of the BH Norm was delayed due to the COVID-19 pandemic. Data users can expect to see a revised BH Norm update for the 2025 survey. The last BH Norm update was completed in 2018.

Scanning across the charts, it is important to observe the factors that differ the most from the Bach Harrison Norm. This is the first step in identifying the levels of risk and protection that are higher or lower than those in other communities. The risk factors that are higher than the Bach Harrison Norm and the protective factors that are lower than the Bach Harrison Norm are factors your community should consider addressing when planning prevention programs.

The Xs represent national mental health data gathered by the Youth Risk Behavior Survey (YRBS). National comparison points are available for grades 10 and 12 on the topic of suicide and depression.

Substance Use

There are two types of use measured on the drug use charts.

State Identified Priority Substance Use measures lifetime and 30-day use rates for alcohol, marijuana, tobacco (including e-cigarettes), prescription narcotics, and overall prescription drug abuse.

Other Substance Use measures lifetime and 30-day use rates for a variety of illicit drugs, including cocaine, heroin, and methamphetamine.

Risky Substance Use-Related Behavior is measured in several different ways: binge drinking (having five or more drinks in a row during the two weeks prior to the survey), use of one-half a pack or more of cigarettes per day, and youth indicating drinking alcohol and driving or reporting riding with a driver who had been drinking alcohol during the past 30 days.

Substance Use Treatment Needs are estimates of youth in need of alcohol and drug treatment as well as an estimate of students that need either alcohol OR drug treatment.

The need for substance use treatment is defined as students who report using alcohol on 10 or more occasions in their lifetime or any drugs in their lifetime and marked at least three of the

following items specific to their drug or alcohol use in the past year:

- Spent more time using than intended
- Neglected some of your usual responsibilities because of use
- Wanted to cut down on use
- Others objected to your use
- Frequently thought about using
- Used alcohol or drugs to relieve feelings such as sadness, anger, or boredom

Students could mark whether these items related to their drug use and/or their alcohol use.

Antisocial Behavior

Antisocial Behavior (ASB) profiles show the percentage of youth who reported antisocial behaviors during the past year, including suspension from school, selling illegal drugs, and attacking another person with the intention of doing them serious harm.

Community and School Climate and Safety

These charts present data related to perceived importance of school, positive school environment, student concerns about their school environment, perceived school safety, and bullying. The section also includes data for perceived reasons of bullying, with data being presented of students reporting being bullied in the past year (a subset of the total survey sample).

Social and Emotional Health

The social and emotional health data show the percentage of youth with depression and mental health treatment needs, student responses to questions about suicide and self-harm, student responses to questions about social isolation, and questions about student attitudes toward the acceptability of seeking mental health treatment and their willingness to do so.

Needs Mental Health Treatment was estimated using the K6 Scale that was developed with support from the National Center for Health Statistics for use in the National Health Interview Survey. The tool screens for psychological distress by asking students

During the past 30 days, how often did you:

- feel nervous?
- feel hopeless?
- feel restless or fidgety?
- feel so depressed that nothing could cheer you up?
- feel that everything was an effort?
- feel worthless?

Answers to each were scored based on responses: None of the time (0 points), A little of the time (1 point), Some of the time (2 points), Most of the time (3 points), All of the time (4 points). Students with a total score of 13 or more points were determined to have high mental health treatment needs. Table 6 also shows the percentage of students with moderate (scoring 7-12 points) and low (scoring 0-6 points) mental health treatment needs.

Depression-Related Indicators are divided into two sections. The first asks about depression in the past year:

During the past 12 months, did you ever feel so sad or hopeless almost every day for two weeks or more in a row that you stopped doing some usual activities?

A second part, the depressive symptoms scale, is also reported. This part is calculated from student responses to the following statements:

- Sometimes I think that life is not worth it.
- At times I think I am no good at all.
- All in all, I am inclined to think that I am a failure.
- In the past year, have you felt depressed or sad MOST days, even if you felt OK sometimes?

These four depressive symptoms questions were scored on a scale of 1 to 4 (NO!, no, yes, YES!). The survey respondents were divided into three groups. The first group was the High Depressive Symptoms group who scored at least a mean of 3.75 on the depressive symptoms. This meant that those individuals marked *Definitely Yes* to all four items or marked *Somewhat Yes* to one item and *Definitely Yes* to three. The second group was the No Depressive Symptoms group who marked *Definitely No* to all four of the items, and the third group was a middle group who comprised the remaining respondents.

Suicide Related Indicators are based on a series of questions about suicide. These questions provide information about suicidal ideation and attempts of suicide (e.g., "During the past 12 months, did you ever seriously consider attempting suicide?" and "During the past 12 months, how many times did you actually attempt suicide?").

Self-Harm question (introduced in 2015) asks about self-destructive behavior other than suicide. Students are considered to have engaged in self-harm if they responded they had done "something to purposefully hurt yourself without wanting to die, such as cutting or burning yourself on purpose" one or more times during the past 12 months.

Attitudes Toward Mental Health Treatment are explored in a series of questions introduced in the 2017 SHARP survey. How often they talked to an adult about "feeling very sad, hopeless, or suicidal," and if so, who they talked with. The final question in this section explores student attitudes toward seeking professional mental health treatment when they are feeling this way.

Substance-Related Behaviors

The charts in this section explore additional data related to substance use. The charts present data regarding student perceptions of parental disapproval, student attitudes toward peer use, and student perceptions of how harmful substance use is. Further, the charts explore student places of alcohol use (data are presented of past-year alcohol users), as well as student sources of obtaining e-cigarette/vaping devices and places of e-cigarette/vape device use (of students who have vaped in the past year).

Physical Health and Safety

This section focuses on student physical health and safety. Charts display the percent of students reporting safe vehicle habits and positive habits related to physical health and activity. New screen time and device use data are also reported in this section.

Risk and Protective Factors

Risk and protective factor scales measure specific aspects of a youth's life experience that predict whether he/she will engage in problem behaviors. The scales, defined in Table 2, are grouped into four domains: community, family, school, and peer/ individual. The risk and protective factor charts show the percentage of students at risk and with protection for each of the scales.

Risk Factor Charts show the percentage of youth who are considered "higher risk" across a variety of risk factor scales.

Protective Factor Charts show the percentage of youth who are considered high in protection across a variety of protective factor scales.

Additional Tables in this Report

Appendices B and C contain additional data for prevention planning and reporting to state and federal agencies.

Perceived Parental Approval and ATOD Use

Appendix B explores the relationship between perceived parental approval and ATOD use. A full explanation of how to interpret these data is available accompanying the tables.

Drug Free Communities

Appendix C contains information relevant to Drug Free Community (DFC) grantees. This table reports the four DFC Core Measures on alcohol, tobacco, marijuana and prescription drugs:


Perception of Risk - The percentage of respondents who report that regular use of the substance has *moderate risk* or *great risk*.

Perception of Parental Disapproval - The percentage of respondents who report their parents would feel regular use of alcohol or any use of cigarettes, e-cigarettes, marijuana, or the misuse of prescription drugs is *wrong* or *very wrong*.

Perception of Peer Disapproval - The percentage of respondents who report their friends would feel regular use of alcohol or any use of cigarettes, marijuana, or misuse of prescription drugs is *wrong* or *very wrong*.

Past 30-Day Use - The percentage surveyed reporting using the substance at least once in the past 30 days.

State-identified priority substance use - Alcohol, marijuana, and prescription drugs White Students 2023 SHARP (PNA) Student Survey




^{*} Since not all students answer all questions, the percentage of students reporting use in the past 30 days may be greater than the percentage reporting lifetime use.

Race 2019 Race 2021 State 2023 ♦ MTF** Race 2023

^{**} National comparison data are available for 8th, 10th, and 12th grade only. Monitoring the Future does not survey 6th graders.

^{† &}quot;Prescription drug abuse" is a combined measure showing the total rate of abuse of any prescription stimulant, prescription sedative, prescription tranquilizer, or prescription narcotic drugs.

How marijuana was used (of past-month marijuana users)* White Students 2023 SHARP (PNA) Student Survey




				During the past 30) days, how did you ι	ıse marijuana?*(Marl	〈 ALL that apply.)					
	l smo	ked it	l ate it (in an edibl or othe		I used a v	vaporizer	l dabl	oed it	l used it in some other way			
Grade	Race 2023	State 2023	Race 2023	State 2023	Race 2023	State 2023	Race 2023	State 2023	Race 2023	State 2023		
8th	73.7	61.3	17.9	18.0	63.4	49.1	38.0	36.8	15.1	21.4		
10th	76.3	77.7	28.5	28.8	43.8	40.8	17.9	25.2	0.0	6.1		
12th	87.2	64.6	19.3	19.0	35.0	52.6	15.6	15.4	2.5	2.3		
All	82.2	68.3	22.3	21.7	40.6	48.7	18.5	20.3	2.8	5.6		

^{*} These data represent youth who had used marijuana in the past month and chose at least one source of consuming it. If this represents fewer than ten students, the sample is considered too small to produce reliable results and the data are removed. If no data are available, state data are displayed instead. The 2023 sample sizes represented in these data are 6th grade: <10 students (State: <30); 8th grade: 19 students (State: 35); 10th grade: 47 students (State: 85); 12th grade: 38 students (State: 76); all grades combined: 104 students (State: 198).

Race 2023 State 2023

State-identified priority substance use - Prescription drugs White Students 2023 SHARP (PNA) Student Survey




^{*} Since not all students answer all questions, the percentage of students reporting use in the past 30 days may be greater than the percentage reporting lifetime use.

Race 2019 Race 2021 Race 2023 State 2023 ♦ MTF**

^{**} National comparison data are available for 8th, 10th, and 12th grade only. Monitoring the Future does not survey 6th graders.


State-identified priority substance use - Cigarettes and vaping products White Students 2023 SHARP (PNA) Student Survey



^{*} Since not all students answer all questions, the percentage of students reporting use in the past 30 days may be greater than the percentage reporting lifetime use

^{**} National comparison data are available for 8th, 10th, and 12th grade only. Monitoring the Future does not survey 6th graders.

^{***} In 2019 and 2021, lifetime and 30-day vaping use questions were about general vape device use. In 2023, the survey was revised to ask about vaping nicotine and vaping marijuana separately. The 2023 vaping data presented here are for any vaped nicotine OR vaped marijuana.




^{*} Since not all students answer all questions, the percentage of students reporting use in the past 30 days may be greater than the percentage reporting lifetime use.

Race 2019 Race 2021 Race 2023 • State 2023 • MTF**

^{**} National comparison data are available for 8th, 10th, and 12th grade only. Monitoring the Future does not survey 6th graders.

Other drugs - Cocaine, methamphetamine, heroin White Students 2023 SHARP (PNA) Student Survey




^{*} Since not all students answer all questions, the percentage of students reporting use in the past 30 days may be greater than the percentage reporting lifetime use.

Race 2019 Race 2021 Race 2023 State 2023 ♦ MTF**

^{**} National comparison data are available for 8th, 10th, and 12th grade only. Monitoring the Future does not survey 6th graders.

Other drugs - Inhalants, hallucinogens, steroids, synthetics White Students 2023 SHARP (PNA) Student Survey




^{*} Since not all students answer all questions, the percentage of students reporting use in the past 30 days may be greater than the percentage reporting lifetime use.

Race 2019 Race 2021 Race 2023 State 2023 ♦ MTF**

^{**} National comparison data are available for 8th, 10th, and 12th grade only. Monitoring the Future does not survey 6th graders.

Risky substance use-related behavior White Students 2023 SHARP (PNA) Student Survey




^{*} Since not all students answer all questions, the percentage of students reporting binge drinking may be greater than the percentage reporting 30-day alcohol use.

Race 2019 Race 2021 Race 2023 State 2023 ▲ BH Norm ♦ MTF**

^{**} National comparison data are available for 8th, 10th, and 12th grade only. Monitoring the Future does not survey 6th graders.


Substance treatment needs White Students 2023 SHARP (PNA) Student Survey

Race 2019



Race 2021 Race 2023

Antisocial behavior White Students 2023 SHARP (PNA) Student Survey




Perceived importance of school White Students 2023 SHARP (PNA) Student Survey


Race 2019

Race 2021

Race 2023



Positive school environment White Students 2023 SHARP (PNA) Student Survey




Student concerns about school environment White Students 2023 SHARP (PNA) Student Survey

Race 2019


Race 2021

Race 2023



Perceived school safety White Students 2023 SHARP (PNA) Student Survey

Race 2019




Race 2021 Race 2023

Bullying White Students 2023 SHARP (PNA) Student Survey


Race 2019

Race 2021

Race 2023



Mental health treatment needs White Students 2023 SHARP (PNA) Student Survey



^{*} Mental health treatment needs are calculated from student responses to specific questions. See text for further explanation.


Race 2019 Race 2021 Race 2023 State 2023

Attitudes toward mental health treatment

		6th			8th				10th				12th				All				
		Race 2019	Race 2021	Race 2023	State 2023																
How often in the last thirty days did you talk to an adult (parent, doctor, counselor, teacher, etc.) about feeling very sad, hopeless, or suicidal? (Answered 1 or more times)		16.5	17.6	22.2	23.5	16.2	18.3	19.3	19.7	22.5	22.2	24.2	23.8	23.0	25.8	25.6	25.1	19.7	21.1	22.9	23.0
	Sample size*	2,386	987	1,463	2,367	2,910	2,162	1,786	2,586	3,146	2,370	1,689	2,245	1,987	1,445	929	1,208	10,42 9	6,964	5,867	8,406
	I felt this way but did not talk to anyone about it	28.1	40.2	39.3	43.9	34.5	46.1	44.8	48.6	31.2	37.1	41.8	43.2	30.8	32.5	34.5	36.3	31.3	38.1	39.8	42.7
	Parent	60.3	45.8	48.2	43.0	46.8	32.5	36.2	32.0	42.4	35.4	39.2	36.1	42.8	40.8	39.8	37.7	46.7	37.9	40.3	37.0
Who, in the last thirty	Friend/Peer	37.7	29.6	35.5	34.1	49.1	41.1	41.7	39.8	55.8	48.8	45.3	43.8	52.1	52.8	50.2	50.1	50.0	45.6	44.3	42.7
days, did you talk to about feeling very sad,	Teacher	15.0	4.8	9.8	9.6	6.6	2.8	4.3	4.5	5.5	3.1	3.5	3.3	6.3	4.6	4.3	4.0	7.7	3.8	5.0	5.1
hopeless, or suicidal?	Doctor	14.1	4.3	7.6	7.5	8.9	5.5	6.9	5.9	7.8	5.8	7.3	6.4	7.5	7.3	6.1	5.3	9.1	6.0	6.9	6.2
(Treated as "Mark all that apply")	Clergy (e.g. Bishop, Priest or Nun, Minister, Pastor)	8.4	2.5	4.0	3.3	6.1	1.4	2.3	1.8	5.1	2.0	2.7	2.4	4.4	3.4	2.8	3.0	5.7	2.4	2.9	2.6
	School Counselor	17.6	10.4	13.7	13.9	11.5	6.6	8.1	7.9	6.8	4.3	3.9	4.4	5.8	3.0	3.2	3.4	9.5	5.2	6.3	6.9
	School Nurse	~	0.4	1.5	1.5	~	0.4	0.5	0.6	~	0.5	0.1	0.3	~	0.1	0.0	0.1	~	0.3	0.4	0.5
	Therapist	20.6	14.9	17.3	17.5	15.1	12.8	15.3	14.1	16.2	14.2	19.7	18.0	14.7	16.6	20.5	19.1	16.3	14.8	18.6	17.3
	Other adult	15.4	6.7	13.6	12.9	10.7	7.3	9.8	9.8	7.7	6.4	8.1	8.3	7.6	5.8	8.0	9.8	9.7	6.5	9.4	10.0
Do you think it's ok to seek help and talk to a professional counselor, therapist,	Yes	91.0	89.0	87.5	84.0	86.7	85.9	85.2	81.0	85.5	84.2	82.0	80.1	87.0	86.8	85.7	83.7	87.5	86.3	85.0	82.2
or doctor if you've been feeling very sad, hopeless, or suicidal?	No	2.5	2.2	2.2	3.4	2.3	1.5	1.7	3.0	2.1	1.6	2.3	2.7	2.0	1.9	2.8	3.1	2.2	1.8	2.3	3.0
Do you think it's ok to seek help and talk to a professional counselor, therapist, or doctor if you've been feeling very sad, hopeless, or suicidal?	I think it's ok for other people to seek help, but not for me to seek help	6.5	8.9	10.3	12.6	11.0	12.6	13.1	16.0	12.4	14.2	15.7	17.2	11.0	11.3	11.5	13.2	10.3	11.9	12.8	14.8

^{*} These data represent youth who chose any answer other than "I have not felt this way in the past 30 days." If this represents fewer than ten students, the sample is considered too small to produce reliable results and the data are removed. Due to changes to this question, 2017 data were unsuitable for comparison and are omitted. See Appendix A for information.

Depression White Students 2023 SHARP (PNA) Student Survey




^{*} Depressive symptoms are calculated from student responses to specific questions. See text for further explanation.

Race 2019 Race 2021 Race 2023 State 2023


^{**} National comparison data for "Felt sad or hopeless..." are available for 10th and 12th grade only.

Suicide indicators and self harm White Students 2023 SHARP (PNA) Student Survey




Social isolation White Students 2023 SHARP (PNA) Student Survey

Race 2019




Race 2021 Race 2023

Social Determinants of Health White Students 2023 SHARP (PNA) Student Survey




Places of alcohol use (of past-year alcohol users)* White Students 2023 SHARP (PNA) Student Survey



^{*}These data represent youth who had drank alcohol in the past month and chose at least one place for consuming it. If this represents fewer than ten students, the sample is considered too small to produce reliable results and the data are removed. If no data are available, state data are displayed instead. The 2023 sample sizes represented in these data are 6th grade; 195 students (State; 427); 8th grade; 489 students (State; 916); 10th grade; 692 students (State; 1,121); 12th grade; 648 students (State; 927); all grades combined; 2,024 students (State; 3,391).

Race 2019 Race 2023 State 2023


Sources of vaping products (of past-month vaping product users)* White Students 2023 SHARP (PNA) Student Survey



^{*} These data represent youth who had used a vaping device in the past month and chose at least one source of obtaining vaping products. If this represents fewer than ten students, the sample is considered too small to produce reliable results and the data are removed. If no data are available, state data are displayed instead. The 2023 sample sizes represented in these data are 6th grade: 29 students (State: 29); 8th grade: 140 students (State: 276); 10th grade: 167 students (State: 265); 12th grade: 127 students (State: 174); all grades combined: 463 students (State: 276); 10th grade: 167 students (State: 276); 12th grade: 127 students (State: 804). Race 2019 Race 2021 Race 2023 State 2023

2023 Prevention Needs Assessment Survey


Sources of cigarettes (of past-month cigarette users)* White Students 2023 SHARP (PNA) Student Survey



^{*}These data represent youth who had smoked cigarettes in the past month and chose at least one source of obtaining them. If this represents fewer than ten students, the sample is considered too small to produce reliable results and the data are removed. If no data are available, state data are displayed instead. The 2023 sample sizes represented in these data are 6th grade: <10 students (State: 10); 8th grade: 23 students (State: 41); 10th grade: 23 students (State: 28); all grades combined: 73 students (State: 120).

Race 2019 Race 2021 Race 2023 State 2023


First use of tobacco (of students who have ever tried tobacco products)* White Students 2023 SHARP (PNA) Student Survey




^{*}These data do not include students who have never used a tobacco product. If this represents fewer than ten students, the sample is considered too small to produce reliable results and the data are removed. If no data are available, state data are displayed instead. The 2023 sample sizes represented in these data are 6th grade: 101 students (State: 227); 8th grade: 336 students (State: 635); 10th grade: 382 students (State: 618); 12th grade: 296 students (State: 442); all grades combined: 1,115 students (State: 1,922).

Race 2019 Race 2021 Race 2023 State 2023


Perception of parental disapproval White Students 2023 SHARP (PNA) Student Survey



Perception of peer disapproval White Students 2023 SHARP (PNA) Student Survey



Perception of risk White Students 2023 SHARP (PNA) Student Survey



Perceived vs. actual ATOD use*

	6th					8t	h			10	th			12	th			Α	II		
		Race 2019	Race 2021	Race 2023	State 2023																
Smoke cigarettes every day	Perceived use	2.9	2.7	2.5	2.7	12.3	11.4	10.8	11.7	18.3	16.2	15.1	16.1	18.2	14.9	13.7	14.3	13.0	11.6	10.9	11.2
0 , ,	Actual use	0.1	0.1	0.0	0.0	0.1	0.1	0.0	0.1	0.4	0.3	0.2	0.2	0.6	0.3	0.3	0.2	0.3	0.2	0.1	0.2
Drank alcohol in past 30 days	Perceived use	3.7	3.5	3.1	3.2	16.9	15.0	12.9	13.9	28.8	26.0	21.9	23.2	31.8	27.6	26.0	26.8	20.4	18.6	16.6	16.7
,	Actual use	0.5	0.5	0.4	0.8	2.2	1.9	1.3	2.0	5.6	3.8	2.8	4.0	9.2	7.1	6.7	7.2	4.3	3.4	2.9	3.4
Used marijuana in past 30	Perceived use	2.1	1.7	1.7	2.1	15.4	12.1	11.8	14.2	29.6	23.6	20.2	22.6	31.8	26.4	23.5	25.4	19.8	16.5	14.9	16.1
days	Actual use	0.2	0.2	0.2	0.6	2.4	1.6	1.7	3.1	7.0	4.8	3.7	5.1	9.2	8.0	7.4	8.6	4.7	3.7	3.4	4.3
Used a vape product	Perceived use	~	~	4.6	5.4	~	~	20.5	23.5	~	~	30.8	33.6	~	~	34.2	35.5	~	~	23.3	24.5
osed a vape product	Actual use	~	~	0.7	1.5	~	~	2.7	4.2	~	~	4.5	6.3	~	~	6.6	7.7	~	~	3.7	4.9

^{*} Students were asked to report the percent of their peers they perceived were using cigarettes, alcohol, and marijuana. The data in this table represent their perceived use amongst similarly aged peers compared to the actual use of those substances by grade.


Student health habits and routines White Students 2023 SHARP (PNA) Student Survey

Race 2019

Race 2021

Race 2023

State 2023




Student physical health and activity levels White Students 2023 SHARP (PNA) Student Survey

Race 2019


Race 2021

Race 2023

State 2023



Safe vehicle habits White Students 2023 SHARP (PNA) Student Survey



Race 2019 Race 2021 Race 2023 • State 2023


Student safety White Students 2023 SHARP (PNA) Student Survey

Race 2019

Race 2021

Race 2023

State 2023



THE RISK AND PROTECTIVE FACTOR MODEL OF PREVENTION

Prevention is a science. The Risk and Protective Factor Model of Prevention is a proven way of reducing substance abuse and its related consequences.

This model is based on the simple premise that to prevent a problem from happening, we need to identify the factors that increase the risk of that problem developing and then find ways to reduce the risks. Just as medical researchers have found risk factors for heart disease such as diets high in fat, lack of exercise, and smoking; a team of researchers at the University of Washington have defined a set of risk factors for youth problem behaviors.

Risk factors are characteristics of school, community, and family environments, and of students and their peer groups known to predict increased likelihood of drug use, delinquency, school dropout, and violent behaviors among youth. For example, children who live in disorganized, crime-ridden neighborhoods are more likely to become involved in crime and drug use than children who live in safe neighborhoods.

The chart on the following page shows the links between the 20 risk factors and six problem behaviors. The check marks indicate where at least two well designed, published research studies have shown a link between the risk factor and the problem behavior.

Protective factors exert a positive influence and buffer against the negative influence of risk, thus reducing the likelihood that adolescents will engage in problem behaviors. Protective factors identified through research include strong bonding to family, school, community and peers, and healthy beliefs and clear standards for behavior. Protective bonding depends on three conditions:

- **Opportunities** for young people to actively contribute
- **Skills** to be able to successfully contribute
- Consistent recognition or reinforcement for their efforts and accomplishments

Bonding confers a protective influence only when there is a positive climate in the schools, families, and neighborhoods. Peers and adults in these schools, families and neighborhoods must communicate healthy values and set clear standards for behavior in order to ensure a protective effect. For example, strong bonds to antisocial peers would not be likely to reinforce positive behavior.

Research on risk and protective factors has important implications for children's academic success, positive youth development, and prevention of health and behavior problems. In order to promote academic success and positive youth development and to prevent problem behaviors, it is necessary to address the factors that predict these outcomes. By measuring risk and protective factors in a population, specific risk factors that are elevated and widespread can be identified and targeted by policies, programs, and actions shown to reduce those risk factors and to promote protective factors.

Each risk and protective factor can be linked to specific types of interventions that have been shown to be effective in either reducing risk(s) or enhancing protection(s). The steps outlined here will help make key decisions regarding allocation of resources, how and when to address specific needs, and which strategies are most effective and known to produce results.


In addition to helping assess current conditions and prioritize areas of greatest need, data from the SHARP Prevention Needs Assessment (PNA) Survey can be a powerful tool in applying for and complying with several federal programs, outlined later in this report, such as the Strategic Prevention Framework process. The survey also gathers valuable data which allows state and local agencies to address other prevention issues related to academic achievement, mental health, gang involvement, health and fitness, and personal safety.

RISK FACTORS FOR ADOLESCENT PROBLEM BEHAVIORS


		Community							Fa	mily		Scho	ol			Pe	er/Indi	vidual		
	Availability of Drugs	Availability of Firearms	Community Laws & Norms Favorable Toward Drug Use, Firearms, & Crime	Media Portrayals of the Behavior	Transitions & Mobility	Low Neighborhood Attachment & Community Disorganization	Extreme Economic Deprivation	Family History of the Problem Behavior	Family Management Problems	Family Conflict	Favorable Parental Attitudes & Involvement in the Problem Behavior	Academic Failure Beginning in Late Elementary School	Lack of Commitment to School	Early & Persistent Antisocial Behavior	Rebelliousness	Gang Involvement	Friends Who Engage in the Problem Behavior	Favorable Attitudes Toward the Problem Behavior	Early Imitation of the Problem Behavior	Constitutional Factors
Substance Abuse	1		1	1	1	√	1	1	√	1	✓	✓	1	1	✓	√	1	✓	1	1
Delinquency		1	/		1	1	1	1	1	1	✓	✓	1	✓	✓	\	1	✓	1	1
Teen Pregnancy							1	1	✓	1		✓	1	1			1	✓	1	
School Drop-Out					1		1	1	1	1		√	1	1	✓		1	√	1	
Violence	1	1	/	1		√	1	1	1	1	√	1	1	1	/	1	1	1	1	1
Depression & Anxiety					1			1	1	√		✓		1						1

2023 Prevention Needs Assessment Survey 43


Risk profile, 6th Grade White Students 2023 SHARP Survey




Risk profile, 8th Grade White Students 2023 SHARP Survey




Risk profile, 10th Grade White Students 2023 SHARP Survey



Risk profile, 12th Grade White Students 2023 SHARP Survey




Risk profile, All Grades White Students 2023 SHARP Survey




Risk profile

	6th							8th					10th			Ī		12th			All				
	Race 2019	Race 2021	Race 2023	State 2023	BH Norm	Race 2019	Race 2021	Race 2023	State 2023	BH Norm	Race 2019	Race 2021	Race 2023	State 2023	BH Norm	Race 2019	Race 2021	Race 2023	State 2023	BH Norm	Race 2019	Race 2021	Race 2023	State 2023	BH Norm
Community Domain																									
Low neighborhood attachment	23.3	24.5	25.1	31.9	42.1	20.7	23.1	24.3	29.9	35.7	27.9	30.0	32.6	36.5	42.8	33.8	36.9	37.8	42.8	49.4	26.4	28.7	30.1	35.1	42.5
Laws & norms favorable to drug use	22.4	28.4	29.7	32.8	35.2	19.1	21.3	21.8	26.2	33.5	16.5	19.4	17.9	20.6	36.0	21.7	22.6	22.8	25.2	44.2	19.8	22.7	22.8	26.2	37.3
Perceived availability of drugs	26.1	25.6	28.1	29.7	35.8	18.9	14.8	14.9	18.2	26.8	22.1	15.0	14.3	16.7	27.4	22.7	15.7	14.3	15.2	32.7	22.4	17.3	17.5	19.9	30.1
Perceived availability of handguns	23.1	22.7	26.7	23.8	22.4	36.5	37.3	35.4	32.6	33.2	24.6	24.0	24.4	23.0	21.8	33.3	30.5	28.2	26.4	26.7	29.4	29.0	28.6	26.5	26.4
Family Domain																									
Poor family management	28.3	30.5	36.5	40.8	44.2	15.3	13.6	16.8	20.1	36.0	14.7	14.1	13.5	15.3	32.0	17.0	15.4	14.2	15.6	35.2	18.7	17.7	19.7	22.9	35.9
Family conflict	28.0	27.9	32.2	34.3	36.9	22.9	23.5	24.0	25.9	32.7	28.6	27.0	26.9	29.0	37.5	27.0	26.2	25.4	27.2	37.5	26.7	26.1	27.0	29.1	36.1
Family history of antisocial behavior	21.0	19.4	19.5	23.9	32.8	15.6	13.8	11.6	14.7	29.5	19.9	16.5	15.1	17.4	32.6	18.0	16.4	16.9	18.5	34.4	18.6	16.4	15.7	18.6	32.3
Parent attitudes favorable to ASB	26.8	31.6	43.2	46.7	36.9	34.2	39.9	45.2	47.4	46.5	35.7	38.6	43.5	45.3	49.6	34.7	38.4	40.8	41.2	49.1	32.9	37.3	43.2	45.2	46.4
Parent attitudes favorable to drug use	4.7	4.9	5.9	7.6	11.4	8.1	8.4	7.8	9.8	22.7	13.7	13.0	12.8	14.6	35.6	12.7	11.6	12.8	14.0	36.8	9.8	9.6	10.0	11.5	28.0
School Domain																									
Academic failure	20.4	22.1	22.1	28.8	27.4	18.6	21.7	22.1	29.7	29.6	21.4	24.1	21.6	28.4	32.3	23.8	26.4	26.2	31.2	33.4	21.0	23.6	23.0	29.5	31.0
Low commitment to school	43.2	45.3	52.6	54.8	47.0	46.9	49.2	54.7	57.1	49.1	44.5	49.1	50.1	50.6	45.9	44.1	44.7	46.6	47.4	47.8	44.7	47.2	50.9	52.5	47.5
Peer-Individual Domain																									
Rebelliousness	16.7	19.8	22.4	26.3	22.0	17.1	17.5	18.6	21.3	26.0	22.2	22.7	22.8	24.7	30.4	23.0	23.7	24.1	25.0	31.7	19.8	21.0	22.0	24.3	28.1
Early initiation of ASB	14.0	14.1	16.1	18.2	24.6	18.2	17.0	19.1	21.9	24.6	20.6	20.7	19.6	21.7	26.3	21.1	19.0	18.6	19.5	25.5	18.4	17.8	18.4	20.4	25.3
Early initiation of drug use	6.0	7.0	7.4	11.5	19.4	8.7	7.6	7.0	10.6	23.3	9.5	8.0	4.9	7.6	24.8	12.4	9.6	6.3	8.5	32.3	9.1	8.1	6.3	9.5	25.7
Attitudes favorable to ASB	37.0	40.5	54.1	57.9	38.8	25.8	30.3	33.5	37.4	29.4	32.9	37.2	39.9	42.0	34.7	34.5	37.8	41.5	40.9	35.3	32.5	36.3	41.9	44.5	34.1
Attitudes favorable to drug use	8.0	9.3	11.8	14.7	17.3	15.5	17.1	16.3	20.1	27.9	21.6	23.5	21.2	23.7	39.6	21.4	22.8	23.2	23.1	41.4	16.6	18.4	18.4	20.4	33.0
Perceived risk of drug use	32.4	38.1	39.0	42.5	50.9	27.5	28.4	28.0	32.1	47.7	33.2	35.0	31.1	35.1	60.2	30.6	31.5	31.5	32.8	58.6	31.0	33.1	32.2	35.6	54.5
Interaction with antisocial peers	17.6	15.2	18.4	21.9	31.7	13.4	9.5	11.6	15.0	23.9	14.4	10.7	10.9	13.3	24.0	14.1	10.6	11.0	11.3	23.5	14.9	11.3	12.8	15.5	25.2
Friend's use of drugs	5.7	5.3	5.7	8.3	14.6	11.8	8.7	8.7	13.0	26.5	11.9	8.0	6.8	8.6	27.7	9.9	7.8	6.3	6.8	28.2	9.9	7.6	6.9	9.2	25.4
Rewards for ASB	19.9	21.2	21.7	23.4	21.6	28.3	25.9	25.9	27.4	30.1	29.8	26.0	24.4	26.1	39.5	28.9	25.4	23.4	24.0	44.1	26.9	24.9	23.9	25.2	35.1
Depressive symptoms	24.0	28.2	33.9	39.0	31.1	31.5	37.0	38.4	41.7	37.4	39.1	46.5	45.8	47.3	43.2	38.3	45.0	45.7	46.2	41.8	33.2	39.5	41.3	43.6	38.7
Gang involvement	1.0	1.0	1.5	2.1	7.7	1.5	0.9	1.0	2.2	7.4	1.6	1.0	0.8	1.2	7.2	1.0	0.7	0.8	1.1	7.9	1.3	0.9	1.0	1.7	7.5


Protective profile, 6th Grade White Students 2023 SHARP Survey




Protective profile, 8th Grade White Students 2023 SHARP Survey




Protective profile, 10th Grade White Students 2023 SHARP Survey



Protective profile, 12th Grade White Students 2023 SHARP Survey



Protective profile, All Grades White Students 2023 SHARP Survey



Protective profile

		6th					8th					10th					12th					All		- 1	
	Race 2019	Race 2021	Race 2023	State 2023	BH Norm	Race 2019	Race 2021	Race 2023	State 2023	BH Norm	Race 2019	Race 2021	Race 2023	State 2023	BH Norm	Race 2019	Race 2021	Race 2023	State 2023	BH Norm	Race 2019	Race 2021	Race 2023	State 2023	BH Norm
Community Domain																									
Rewards for prosocial involvement	66.1	63.3	62.3	54.2	41.8	68.7	64.9	64.4	55.1	45.1	65.3	61.4	62.6	54.0	39.7	63.1	61.2	60.8	53.5	38.9	65.8	62.7	62.5	54.2	41.4
Family Domain																									
Family attachment	80.3	77.5	76.7	70.0	63.5	77.9	73.9	75.0	69.7	59.9	76.9	73.3	76.3	72.7	61.4	75.9	72.9	75.0	71.4	59.0	77.7	74.2	75.7	71.0	60.6
Opportunities for prosocial involvement	80.1	77.1	75.2	69.5	57.2	81.0	78.2	77.1	71.9	65.9	75.4	72.3	73.8	70.6	60.6	74.2	72.8	72.5	69.0	58.3	77.6	75.0	74.6	70.3	60.7
Rewards for prosocial involvement	69.4	70.8	68.7	63.2	57.7	68.2	64.6	65.2	60.1	52.5	69.9	68.1	70.5	67.1	57.9	68.2	67.6	69.3	65.7	54.6	68.9	67.6	68.5	64.0	55.6
School Domain																									
Opportunities for prosocial involvement	70.4	68.6	70.4	68.7	58.8	75.2	76.3	73.7	73.0	68.5	77.8	77.2	78.5	78.2	66.9	76.9	77.6	78.3	77.7	67.5	75.1	75.1	75.4	74.4	66.2
Rewards for prosocial involvement	67.1	70.9	66.5	64.6	54.6	56.7	61.7	56.3	55.8	54.9	63.9	65.0	66.3	66.1	60.8	53.8	57.7	55.8	56.5	49.4	60.5	63.7	61.2	60.8	55.1
Peer-Individual Domain																									
Belief in the moral order	76.2	71.7	68.2	61.9	62.2	80.4	78.1	74.8	69.7	71.8	72.3	67.0	66.5	64.1	60.6	71.7	68.1	64.2	63.8	58.8	75.1	71.2	68.4	64.9	63.5
Interaction with prosocial peers	51.1	39.0	37.5	31.4	52.3	65.1	56.0	56.7	49.1	54.8	71.9	64.6	69.2	61.3	53.6	69.6	64.9	65.7	59.8	47.6	64.6	57.0	58.0	50.3	52.1
Prosocial involvement	54.9	44.2	48.6	42.7	54.4	61.3	51.8	54.2	48.9	55.4	65.5	58.4	60.5	55.3	57.3	64.6	58.9	59.4	54.5	54.5	61.5	53.6	56.0	50.4	55.5
Rewards for prosocial involvement	58.5	54.3	47.9	44.4	53.9	58.7	58.1	53.2	50.9	54.4	72.3	66.2	66.7	64.4	61.8	74.6	69.7	71.0	69.3	63.3	66.2	62.6	60.2	57.1	58.9

SCHOOL AND COMMUNITY IMPROVEMENT USING SURVEY DATA

What are the numbers telling you?

Review the charts and data tables presented in this report. Note your findings as you discuss the following questions.

- Which 3-5 risk factors appear to be higher than you would want when compared to the Bach Harrison Norm?
- Which 3-5 protective factors appear to be lower than you would want when compared to the Bach Harrison Norm?
- Which levels of 30-day drug use are increasing and/or unacceptably high?
 - Which substances are your students using the most?
 - At which grades do you see unacceptable usage levels?
- Which antisocial behaviors are increasing and/or unacceptably high?
 - Which behaviors are your students exhibiting the most?
 - · At which grades do you see unacceptable behavior levels?

How to identify high priority problem areas.

Once you have familiarized yourself with the data, you can begin to identify priorities.

- **Look across the charts** for items that stand out as either much higher or much lower than the others.
- Compare your data with statewide, and/or national data.
 Differences of 5% between local and other data are probably significant.
- Prioritize problems for your area according to the issues

you've identified. Which can be realistically addressed with the funding available to your community? Which problems fit best with the prevention resources at hand?

 Determine the standards and values held within your community. For example: Is it acceptable in your community for a percentage of high school students to drink alcohol regularly as long as that percentage is lower than the overall state rate?

Use these data for planning.

Once priorities are established, use data to guide your prevention efforts.

- Substance use and antisocial behavior data are excellent tools to raise awareness about the problems and promote dialogue.
- Risk and protective factor data can be used to identify exactly where the community needs to take action.
- Additional survey data on academic achievement, mental health and suicide, health and fitness, gang involvement, and other areas can be used to broaden your prevention approach. Find ways to share these data with other prevention planners in your community.
- **Promising approaches** for any prevention goal are available for through resources listed on the last pages of this report. These contacts are a great resource for information about programs that have been proven effective in addressing the risk factors that are high in your community, and improving the protective factors that are low.

	Sample notes	Priority rate 1	Priority rate 2	Priority rate 3
Risk factors	8th grade, Risk of Drugs seems low @14% (8% > BH Norm.)			
Protective factors	10th grade School rewards for prosocial involvement down 7% from 2 yrs ago			
Substance abuse	8th grade 30-day Vape @7% (3% above state av.)			

BUILDING A STRATEGIC PREVENTION FRAMEWORK

The Prevention Needs Assessment (PNA) Survey is an important data source for communities in creating planned, data-driven, effective, and sustainable prevention programs. The State of Utah endorses two models for guiding prevention work at the community, regional, or State level – the Communities That Care (CTC) Model and the Substance Abuse and Mental Health Services Administration (SAMHSA) Center for Substance Abuse Prevention (CSAP) Strategic Prevention Framework (SPF). Communities in the State of Utah are encouraged to follow the CTC Model, a tested and effective model to guide communities through a process of community organization and mobilization. The second model for prevention planning, the SPF Model, guides states and communities through a five-step process to increase effectiveness of prevention efforts. The following websites provide additional information about these prevention models: http://www.communitiesthatcare.net and

Following are the five steps involved in the SPF model. For training in the SPF or the CTC, contact your local prevention coordinator (http://dsamh.utah.gov/prevention/).

http://www.samhsa.gov/spf.

Assessment: Profile Population Needs, Resources, and Readiness to Address the Problems and Gaps in Service Delivery. The SPF begins with an assessment of the needs in the community that is based on data. The Utah State Epidemiological Outcomes Workgroup (SEOW) has compiled data from several sources to aid in the needs assessment process. One of the primary sources of needs assessment data is this Prevention Needs Assessment Survey (PNA). While planning prevention services, communities are urged to collect and use multiple data sources, including archival and social indicators, assessment of existing resources, key informant interviews, and community readiness. The PNA results presented in this profile report will help you to identify needs for prevention services. PNA data include adolescent substance use,

Evaluation

Sustainability and Cultural Competence

Implementation

Planning

anti-social behavior, and many of the risk and protective factors that predict adolescent problem behaviors.

Capacity: Mobilize and/or Build Capacity to Address Needs. Engagement of key stakeholders at the state and community levels is critical to plan and implement successful prevention activities that will be sustained over time. Some of the key tasks to mobilize the state and communities are to work with leaders and stakeholders to build coalitions, provide training, leverage resources, and help sustain prevention activities.

Planning: Develop a Comprehensive Strategic Plan. States and communities should develop a strategic plan that articulates not only a vision for the prevention activities, but also strategies for organizing and implementing prevention efforts. The strategic

plan should be based on the assessments conducted during Step 1. The Plan should address the priority needs, build on identified resources/strengths, set measurable objectives, and identify how progress will be monitored. Plans should be adjusted with ongoing needs assessment and monitoring activities.

Implementation: Implement Evidence-based Prevention Programs and Infrastructure Development Activities. By measuring and identifying the risk factors and other causal factors that contribute to the targeted problems specified in your strategic plan, programs can be implemented that will reduce the prioritized substance abuse problems. After completing Steps 1, 2, and 3, communities will be able to choose prevention strategies that have been shown to be effective, are appropriate for the population served, can be implemented with fidelity, are culturally appropriate, and can be sustained over time. SAHMSA's National Registry of Evidence-based Programs and Practices (located at http://www.nrepp.samhsa.gov) is a searchable online registry of mental health and substance abuse interventions that have been reviewed and rated by independent reviewers. This resource can help identify scientifically based approaches to preventing and treating mental and/or substance use disorders that can be readily disseminated to the field.

Evaluation: Monitor Process, Evaluate Effectiveness, Sustain Effective Programs/Activities, and Improve or Replace Those That Fail. Finally, ongoing monitoring and evaluation are essential to determine if the desired outcomes are achieved, assess service delivery quality, identify successes, encourage needed improvement, and promote sustainability of effective policies, programs, and practices. The PNA allows communities to monitor levels of ATOD use, antisocial behavior, risk, and protection.

Sustainability and Cultural Competence are at the core of the SPF model, indicating the key role they play in each of the five

elements. Incorporating principles of cultural competence and sustainability throughout assessment, capacity appraisal, planning, implementation and evaluation helps ensure successful, long lasting prevention programs.

Sustainability is accomplished by utilizing a comprehensive approach. By building adaptive and flexible programs around a variety of resources, funding and organizations, states and communities will build sustainable programs and achieve sustainable outcomes. A strategic plan that dynamically responds to changing issues, data, priorities, and resources is more likely to achieve long term results.

Sharing information gathered during the evaluation stage with key stakeholders, forging partnerships and encouraging creative collaboration all enhance sustainability.

Cultural Competence: Planners need to recognize the needs, styles, values and beliefs of the recipients of prevention efforts. Culturally competent prevention strategies use interventions, evaluations and communication strategies appropriate to their intended community. Cultural issues reflect a range of influences and are not just a matter of ethnic or racial identity. Learning to communicate with audiences from diverse geographic, cultural, economic, social, and linguistic backgrounds can increase program efficacy and ensure sustainable results.

Whether enlisting extended family networks as a prevention resource for single parent households, or ensuring there are resources available to bridge language gaps, cultural competency will help you recognize differences in prevention needs and tailor prevention approaches accordingly.

A one-size-fits-all program is less effective than a program that works with knowledgeable people from the community to develop focused interventions, communication, and support and draws on community-based values and traditions.

APPENDIX A: CHANGES BETWEEN PNA ADMINISTRATIONS

As new issues come to the forefront and new prevention modalities are implemented, the SHARP PNA survey evolves to reflect these concerns.

Weighting procedures for 2023

The weighting procedure used for the 2023 SHARP is the same procedure used since 2015. It starts with a school-level weighting procedure. At the district level and above, Bach Harrison analysts apply a raking ratio estimation, which is a method for adjusting the sampling weights of the sample data based on known population characteristics. This helps ensure that the survey sample reflects the total population of Utah students on grade, gender, and race/ethnicity. For more detailed information on the weighting procedure consult the Comprehensive State Report (to be released in late fall 2023).

New Items for 2023

A new question was added to assess social determinants of health.

This past year, did you experience any of the following? (Mark ALL that apply.)

- One or more people living in my home lost their job
- I had to move or change homes in the past year
- Skipped one or more meals because my family didn't have enough money to buy food
- I had difficulty keeping up with schoolwork because I didn't have access to a reliable computer or internet service
- I did not have a quiet place at home to study
- None of these

Additional items added to the 2023 survey were as follows:

During the past year, did you drink alcohol at any of the following places? (Mark the number of times for each.)

During the past 30 days, how did you use marijuana? (Mark ALL that apply.)

Complete response options for new questions can be found in Appendix F: Item Dictionary.

Adjustments to Response Options

The vape products/types question was segmented into two questions:

- vape products containing nicotine (such as e-cigarettes, vape pens, mods, or pod vapes like JUUL or Puff Bars)
- vape products containing marijuana

For the lifetime and 30-day tobacco questions, a new category was added to both:

 nicotine lozenges (small hard or soft tablets containing nicotine that slowly dissolve in the mouth such as Velo or Rogue)?

For the question regarding perceived sibling substance use and the question regarding perceived peer use of substances, the question was included to ask about perceived use of:

 a vape product (e-cigarettes, vape pens, mods, or pod vapes like JUUL or Puff Bars).

Other Survey Removals and Changes

The following questions were removed from the 2023 SHARP PNA:

If you used alcohol the past year (12 months), how did you get it? (Mark all that apply.) (This question has been shifted to an every-other-year administration schedule.)

If you used a vape product such as e-cigarettes, vape pens, mods, or pod vapes like JUUL or Puff Bars in the past 12 months, what did you put in it? (Mark ALL that apply.)

If you used a vape product (e-cigarettes, vape pens, or mods, or pod vapes like JUUL or Puff Bars) in the past 30 days, where did you use it? (Mark ALL that apply.)

This past year, many youth and families in Utah were affected by the Coronavirus (also known as COVID-19). Did you experience any of the following due to the coronavirus or coronavirus symptoms? (Select one or more responses).

How much do you agree or disagree with the following statement: "My learning improved when my classes were taught online due to COVID-19?"

How much do you agree or disagree with the following statements about LEARNING ONLINE:

To review past changes to the SHARP PNA (e.g. differences between and additions to the 2017, 2019 and 2021 surveys), please contact Bach Harrison.

APPENDIX B: SUBSTANCE USE AND PERCEIVED PARENTAL ACCEPTABILITY

Even a small amount of perceived parental acceptability can lead to substance use.

When parents have favorable attitudes toward drugs, they influence the attitudes and behavior of their children. For example, parental approval of moderate drinking, even under parental supervision, substantially increases the risk of the young person using alcohol. Further, in families where parents involve children in their own drug or alcohol behavior, for example, asking the child to light the parent's cigarette or to get the parent a beer, there is an increased likelihood that their children will become drug users in adolescence.

In the 2023 SHARP PNA Survey, students were asked how wrong their parents felt it was to use alcohol, marijuana, cigarettes, or prescription drugs not prescribed to them. The table below displays lifetime and past 30 days use rates in relation to parents' acceptance of alcohol, marijuana, cigarette, or prescription drug abuse.

In 2023, 90.6% of Utah students indicated that their parents felt it was "Very wrong" for them to use alcohol. The data show that, of those students, relatively few (8.0% lifetime, 1.9% 30-day) actually used alcohol. In contrast, of the 6.8% who marked that their parents agree with use somewhat (i.e. the parent only believes that it is "Wrong," not "Very Wrong"), 35.4% of these students indicated lifetime alcohol use and 12.0% of these students indicated 30-day alcohol use. Similar findings can be observed regarding marijuana, cigarette, and prescription drug use.

The table below illustrates how even a small amount of perceived parental acceptability can lead to substance use. These results make a strong argument for the importance of parents having strong and clear standards and rules when it comes to ATOD use.

How wrong do your parents feel it would be for YOU to:	Student ha	s used:
drink beer, wine, or hard liquor regularly?	Alcohol (lifetime use)	Alcohol (past 30 days)
Very Wrong	8.0%	1.9%
Wrong	35.4%	12.0%
A Little Bit Wrong	58.6%	30.9%
Not Wrong At All	59.1%	38.9%
smoke marijuana?	Marijuana (lifetime use)	Marijuana (past 30 days)
Very Wrong	6.4%	2.6%
Wrong	28.7%	14.3%
A Little Bit Wrong	56.0%	33.4%
Not Wrong At All	64.5%	45.6%
smoke cigarettes?	Cigarettes (lifetime use)	Cigarettes (past 30 days)
Very Wrong	4.1%	0.5%
Wrong	16.5%	2.4%
A Little Bit Wrong	36.8%	15.5%
Not Wrong At All	55.9%	18.1%
use prescription drugs not prescribed to you?	Prescription drugs (lifetime use)	Prescription drugs (past 30 days)
Very Wrong	4.2%	1.2%
Wrong	15.1%	4.3%
A Little Bit Wrong	36.7%	19.6%
Not Wrong At All	27.8%	14.4%

APPENDIX C: DRUG FREE COMMUNITIES DATA

			6	th	8t	th	10	th	12	th.	Ma	Male		nale
Core Measure	Definition	Substance	Percent	Sample										
	take one or two drinks of an alcoholic beverage (beer, wine, liquor) nearly every day	Regular drinking	78.9	11,070	83.6	11,281	87.0	8,565	83.5	4,249	81.8	17,019	85.2	17,829
Perception of Risk*	take five or more drinks of an alcoholic beverage once or twice a week	Binge drinking	80.5	11,135	85.7	11,320	88.9	8,583	86.1	4,266	84.3	17,091	86.8	17,894
(People are at Moderate or Great Risk of	smoke one or more packs of cigarettes per day	Tobacco	84.1	11,175	87.5	11,328	89.5	8,582	87.5	4,263	86.7	17,093	88.0	17,934
harming themselves if they)	smoke marijuana regularly	Marijuana	77.8	10,777	80.5	11,267	78.6	8,552	68.0	4,254	73.3	16,889	79.2	17,646
ii dicy)	use prescription drugs that are not prescribed to them	Prescription drugs	83.6	10,996	87.7	11,248	90.5	8,550	88.3	4,256	86.4	16,955	89.1	17,779
	use vape products such as e-Cigarettes, vape pens, mods, or pod vapes like JUUL or Puff Bars?	e-Cigarettes/va ping	85.4	10,889	88.8	11,233	90.2	8,536	87.5	4,242	86.9	16,892	89.4	17,694
	have one or two drinks of an alcoholic beverage nearly every day	Alcohol	99.5	10,964	99.4	11,086	99.2	8,313	98.6	4,131	99.0	16,582	99.3	17,598
Perception of	smoke cigarettes	Tobacco	99.6	11,003	99.4	11,116	99.3	8,327	99.1	4,143	99.2	16,633	99.5	17,642
Parental Disapproval* (Parents feel it would	smoke marijuana	Marijuana	99.6	10,972	98.7	11,086	96.9	8,298	94.2	4,134	97.0	16,584	97.5	17,592
be Wrong or Very Wrong to)	use prescription drugs not prescribed to you	Prescription drugs	99.5	10,942	99.0	11,086	98.6	8,309	98.8	4,130	98.8	16,570	99.1	17,583
	use vape products such as e-Cigarettes, vape pens, mods, or pod vapes like JUUL or Puff Bars?	e-Cigarettes/va ping	99.7	10,867	99.2	10,994	98.5	8,262	97.6	4,114	98.8	16,448	98.6	17,479
Perception of	have one or two drinks of an alcoholic beverage nearly every day	Alcohol	98.3	11,003	96.7	11,150	95.2	8,362	92.5	4,160	95.4	16,701	95.8	17,665
Peer Disapproval* (Friends feel it would	smoke tobacco	Tobacco	98.8	10,992	97.3	11,147	95.6	8,361	93.7	4,157	95.9	16,696	96.7	17,653
be Wrong or Very Wrong to)	smoke marijuana	Marijuana	98.8	10,950	95.4	11,127	90.0	8,348	82.9	4,157	91.1	16,663	91.8	17,612
, , , , , , , , , , , , , , , , , , ,	use prescription drugs not prescribed to you	Prescription drugs	98.5	10,966	97.3	11,126	96.4	8,350	96.1	4,155	96.8	16,670	97.3	17,619
	had beer, wine, or hard liquor	Alcohol	0.4	11,153	1.3	11,267	2.8	8,486	6.7	4,223	2.5	16,953	3.4	17,855
Past 30-Day Use*	smoked cigarettes	Tobacco	0.2	11,079	0.3	11,192	0.7	8,403	1.3	4,179	0.7	16,783	0.6	17,750
(at least one use in the past 30 days)	used marijuana	Marijuana	0.2	11,135	1.7	11,322	3.7	8,534	7.4	4,236	2.8	17,005	4.0	17,901
	combined results of prescription stimulant/sedative/ narcotics questions	Prescription drugs	1.7	11,124	1.4	11,329	1.3	8,553	1.0	4,252	1.1	17,014	1.6	17,921

^{*} For Past 30-Day Use, Perception of Risk, and Perception of Parental/Peer Disapproval, the "Sample" column represents the sample size - the number of people who answered the question and whose responses were used to determine the percentage. The "Percent" column represents the percentage of youth in the sample answering the question as specified in the definition.

2023 Prevention Needs Assessment Survey

APPENDIX D: CONTACTS FOR PREVENTION

National Contacts

National Institute on Alcohol Abuse and Alcoholism

https://www.niaaa.nih.gov

National Clearinghouse for Alcohol and Drug Information

https://store.samhsa.gov

The National Institute on Drug Abuse (NIDA) Drugs of Abuse Information Clearinghouse

https://www.drugabuse.gov/drugs-abuse

Center for Substance Abuse Prevention

https://www.samhsa.gov/prevention

Monitoring the Future

http://monitoringthefuture.org

National Survey on Drug Use and Health

https://nsduhweb.rti.org/respweb/homepage.cfm

State Contacts

Utah Department of Health and Human Services

195 N. 1950 W. Salt Lake City, UT 84116 https://dhhs.utah.gov

Rob Timmerman Program Manager 288 N. 1460 W. Salt Lake City, UT 84116 385-228-5034 rtimerman@utah.gov

David Watkins Program Administrator 288 N. 1460 W. Salt Lake City, UT 84116 385-226-6125 dwatkins@utah.gov Heidi Peterson Dutson Prevention Administrator and Regional Director 288 N. 1460 W. Salt Lake City, UT 84116 801-842-2674 hdutson@utah.gov

Elise Hardy Program Administrator 288 N. 1460 W. Salt Lake City, UT 84116 385-315-7836 elisehardy@utah.gov

Janae Duncan
Director
Division of Population Health
288 N. 1460 W.
Salt Lake City, UT 84116
385-280-0312
janaeduncan@utah.gov

Amanda Smith
Deputy State Epidemiologist
288 N. 1460 W.
Salt Lake City, UT 84116
385-454-5071
arsmith@utah.gov

Jenny Johnson Assistant Director of Communications 195 North 1950 West Salt Lake City, UT 84116 385-290-7826 Jennyjohnson@utah.gov

Mercedes Rodriguez
Health Program Coordinator
288 N. 1460 W.
Salt Lake City, UT 84116
801-538-6159
myrodriguez@utah.gov

Ryan Carrier Research Consultant 288 N. 1460 W. Salt Lake City, UT 84116 801-889-8710 rcarrier@utah.gov

Eric Tadehara
Assistant Director
Office of Substance Use and Mental Health
288 N. 1460 W.
Salt Lake City, UT 84116
801-699-0895
erictadehara@utah.gov

Anna Fondario
Director
Office of Health Promotion and Prevention
288 N. 1460 W.
Salt Lake City, UT 84116
385-258-8537
afondario@utah.gov

Sarah Hodson
Deputy Director
Office of Health Promotion and Prevention
288 N. 1460 W.
Salt Lake City, UT 84116
385-256-4461
arsmith@utah.gov

Claudia Bohner Epidemiologist 288 N. 1460 W. Salt Lake City, UT 84116 801-538-9274 cbohner@utah.gov

Wei Beadles Epidemiologist 288 N. 1460 W. Salt Lake City, UT 84116 385-266-4529 whou@utah.gov

Regional Directors

Salt Lake, Summit, Utah, and Wasatch Counties Caryn Coltrin South Redwood Health Center 7971 South Redwood Road West Jordan, UT 84088 385-486-5320

Beaver, Garfield, Iron, Juab, Kane, Millard, Piute, Sanpete, Sevier, and Washington Counties

Logan Reid Southwest Prevention 474 West 200 North St. George, UT 84780 435-867-7622 Ireid@sbhcutah.org

Box Elder, Cache, Davis, Morgan, Rich, and Weber Counties

Jess Brewer Weber Human Services 237 26th Street Ogden, UT 84401 435-799-1167 jessb@weberhs.org

Carbon, Emery, Grand, and San Juan Counties

David Watkins 288 N. 1460 W Salt Lake City, UT 84116 385-226-6125 dwatkins@utah.gov

Duchesne, Tooele, and Uintah Counties

Heidi Peterson Dutson 288 N. 1460 W. Salt Lake City, UT 84116 801-842-2674

hdutson@utah.gov

Local Substance Abuse Authority/ County Level Providers

See https://sumh.utah.gov/ for contact information for prevention efforts in your neighborhood.

Bear River

Annie Parker Bear River Health Department 655 East 1300 North Logan, UT 84341 435-792-6509 aparker@brhd.org

Central

Elizabeth Hinckley Central Utah Counseling Center 390 West 100 North Ephraim, UT 84627 435-896-8236 elizabethh@cucc.us

Davis

Amelia Williams
Davis Behavioral Health
934 South Main Street
Layton, UT 84041
801-497-1474
ameliaw@dbhutah.org

Four Corners

Taylor Passarella
Four Corners Community Behavioral Health
105 W 100 N
Price, Utah 84501
435-613-2177
tpassarella@fourcorners.ws

Northeastern

Catherine Jurado Northeastern Counseling Center 285 West 800 South Roosevelt, UT 84066 435-725-6334 catherinei@nccutah.org

Salt Lake

Alysa Stuart
Salt Lake County South Redwood Public Health
Center
7971 S 1825 W
West Jordan, UT 84088
385-226-2639
astuart@slco.org

San Juan

Alyn Mitchell San Juan Counseling 735 South 200 West Suite 1 Blanding, UT 84511 435-485-1020 amitchell@sanjuancc.org

Southwest

Logan Reid Southwest Prevention 474 West 200 North St. George, UT 84780 435-867-7622 Ireid@sbhcutah.org

Summit

Pamella Bello-Straus Summit County Health Department 650 Round Valley Dr. Park City, UT 84060 435-333-1551 pbello@summitcounty.org

Tooele

Peter Clegg Tooele County Health Department 151 North Main Street Tooele, UT 84074 435-277-2363 pclegg@tooelehealth.org

Utah County

Heather Lewis
Utah County Health Department
151 South University Avenue, Suite 2204
Provo, UT 84601
801-851-7188
heatherl@utahcounty.gov

Wasatch

Trudy Brereton Wasatch Mental Health 55 South 500 East Heber, UT 84032 435-657-3228 trudyb@wasatch.org

Weber

Jason Skinner Weber Human Services 237 26th Street Ogden, UT 84401 801-625-3679 jasons@weberhs.org

This Report Was Prepared for the State of Utah by Bach Harrison LLC

http://www.bach-harrison.com R. Steven Harrison, Ph.D. Mary VanLeeuwen Johnstun, M.A. R. Paris Bach-Harrison, B.F.A. G. Apollo Bach-Harrison, B.F.A.

APPENDIX E: RISK AND PROTECTIVE SCALE DEFINITIONS

An explanation of the scales that measure the risk and protective factors shown in the profiles

Community Domain Risk Factors

Low Neighborhood Attachment – Low neighborhood bonding is related to higher levels of juvenile crime and drug selling.

Laws and Norms Favorable Toward Drug Use – Research has shown that legal restrictions on alcohol and tobacco use, such as raising the legal drinking age, restricting smoking in public places, and increased taxation have been followed by decreases in consumption. Moreover, national surveys of high school seniors have shown that shifts in normative attitudes toward drug use have preceded changes in prevalence of use.

Perceived Availability of Drugs and Handguns – The availability of cigarettes, alcohol, marijuana, and other illegal drugs has been related to the use of these substances by adolescents. The availability of handguns is also related to a higher risk of crime and substance use by adolescents.

Community Domain Protective Factors

Rewards for Prosocial Involvement – Rewards for positive participation in activities helps youth bond to the community, thus lowering their risk for substance use.

Family Domain Risk Factors

Poor Family Management – Parents' use of inconsistent and/ or unusually harsh or severe punishment with their children places them at higher risk for substance use and other problem behaviors. Also, parents' failure to provide clear expectations and to monitor their children's behavior makes it more likely that they will engage in drug abuse whether or not there are family drug problems.

Family Conflict – Children raised in families high in conflict, whether or not the child is directly involved in the conflict, appear at risk for both delinquency and drug use.

Family History of Antisocial Behavior – When children are raised in a family with a history of problem behaviors (e.g., violence or ATOD use), the children are more likely to engage in these behaviors.

Parental Attitudes Favorable Toward Antisocial Behavior & Drugs – In families where parents use illegal drugs, are heavy users of alcohol, or are tolerant of children's use, children are more likely to become drug abusers during adolescence. The risk is further increased if parents involve children in their own drug (or alcohol) using behavior, for example, asking the child to light the parent's cigarette or get the parent a beer from the refrigerator.

Family Domain Protective Factors

Family Attachment – Young people who feel that they are a valued part of their family are less likely to engage in substance use and other problem behaviors.

Opportunities for Prosocial Involvement – Young people who are exposed to more opportunities to participate meaningfully in the responsibilities and activities of the family are less likely to engage in drug use and other problem behaviors.

Rewards for Prosocial Involvement – When parents, siblings, and other family members praise, encourage, and attend to things done well by their child, children are less likely to engage in substance use and problem behaviors.

School Domain Risk Factors

Academic Failure – Beginning in late elementary (grades 4-6) academic failure increases the risk of both drug abuse and delinquency. It appears that the experience of failure itself, for whatever reasons, increases the risk of problem behaviors.

Low Commitment to School – Surveys of high school seniors have shown that the use of drugs is significantly lower among students who expect to attend college than among those who do not. Factors such as liking school, spending time on homework, and perceiving the coursework as relevant are also negatively related to drug use.

School Domain Protective Factors

Opportunities for Prosocial Involvement – When young people are given more opportunities to participate meaningfully in important activities at school, they are less likely to engage in drug use and other problem behaviors.

Rewards for Prosocial Involvement – When young people are recognized and rewarded for their contributions at school, they are less likely to be involved in substance use and other problem behaviors.

Peer-Individual Risk Factors

Rebelliousness – Young people who do not feel part of society, are not bound by rules, don't believe in trying to be successful or responsible, or who take an active rebellious stance toward society, are at higher risk of abusing drugs. In addition, high tolerance for deviance, a strong need for independence and normlessness have all been linked with drug use.

Early Initiation of Antisocial Behavior and Drug Use – Early onset of drug use predicts misuse of drugs. The earlier the onset of any drug use, the greater the involvement in other drug use and the greater frequency of use. Onset of drug use prior to the age of 15 is a consistent predictor of drug abuse, and a later age of onset of drug use has been shown to predict lower drug involvement and a greater probability of discontinuation of use.

Attitudes Favorable Toward Antisocial Behavior and Drug Use – During the elementary school years, most children express anti-drug, anti-crime, and pro-social attitudes and have difficulty imagining why people use drugs or engage in antisocial behaviors. However, in middle school, as more youth are exposed to others who use drugs and engage in antisocial behavior, their attitudes often shift toward greater acceptance of these behaviors.

Youth who express positive attitudes toward drug use and anti-

social behavior are more likely to engage in a variety of problem

behaviors, including drug use.

Perceived Risk of Drug Use – Young people who do not perceive drug use to be risky are far more likely to engage in drug use.

Interaction with Antisocial Peers – Young people who associate with peers who engage in problem behaviors are at higher risk for engaging in antisocial behavior themselves.

Friends' Use of Drugs – Young people who associate with peers who engage in alcohol or substance abuse are much more likely to engage in the same behavior. Peer drug use has consistently been found to be among the strongest predictors of substance use among youth. Even when young people come from well-managed families and do not experience other risk factors, spending time with friends who use drugs greatly increases the risk of that problem developing.

Rewards for Antisocial Behavior – Young people who receive rewards for their antisocial behavior are at higher risk for engaging further in antisocial behavior and substance use.

Depressive Symptoms – Young people who are depressed are overrepresented in the criminal justice system and are more likely to use drugs. Survey research and other studies have shown a link between depression and other youth problem behaviors.

Gang Involvement – Youth who belong to gangs are more at risk for antisocial behavior and drug use.

Peer-Individual Protective Factors

Belief in the Moral Order – Young people who have a belief in what is "right" or "wrong" are less likely to use drugs.

Interaction with Prosocial Peers – Young people who associate with peers who engage in prosocial behavior are more protected from engaging in antisocial behavior and substance use.

Prosocial Involvement – Participation in positive school and community activities helps provide protection for youth.

Rewards for Prosocial Involvement – Young people who are rewarded for working hard in school and the community are less likely to engage in problem behavior.

APPENDIX F: ITEM DICTIONARY

DEMOGRAPHICS

How old are you? 10 or younger

11

12

13

14

15 16

17

18

19 or older

What grade are you in? 6th

7th 8th

9th 10th 11th

12th

What is your race? (Select one or more)

American Indian or Alaska Native

Asian

Black or African American

Hispanic or Latino

Native Hawaiian or Other Pacific

Islander White

Are you: Male

(Revised in 2019, then reverted to the 2017 question format in 2021. 2017 text included response options for transgender and other.)

Female

is asking

Some people describe themselves as transgender when their sex at birth does not match the way they think or feel about their gender. Are you transgender?

vhen No, I am not transgender

or I am not sure if I am transgender
I do not know what this question

(New for 2021. Omitted in 6th grade survey.)

Yes, I am transgender

Which of the following best describe you? Heterosexual (straight)

Gay or lesbian (Omitted on 6th grade survey.)

Bisexual Not sure

Think of where you live most of the time. Which of the following people live there with you? (Mark all that apply)

See guestionnaire for complete

list of family members

Think of the adults you live with. What is the highest level of schooling any of them completed?

See questionnaire for complete list of school completion

categories

COMMUNITY: Low neighborhood Attachment

If I had to move, I would miss the neighborhood I now **Definitely No**

live in. Somewhat No Somewhat Yes

Definitely Yes

I like my neighborhood. **Definitely No**

> Somewhat No Somewhat Yes **Definitely Yes**

I'd like to get out of my neighborhood. **Definitely No**

> Somewhat No Somewhat Yes **Definitely Yes**

COMMUNITY: Laws and Norms Favorable to Drug Use

How wrong would most adults in your neighborhood think it was for kids your age:

To use marijuana. Very Wrong

Wrong

A little bit wrong Not wrong at all

to drink alcohol. Very Wrong

Wrong

A little bit wrong Not wrong at all

to smoke cigarettes. Very Wrong

Wrong

A little bit wrong Not wrong at all

If a kid smoked marijuana in your neighborhood would

he or she be caught by the police?

Definitely No Somewhat No Somewhat Yes Definitely Yes

If a kid drank some beer, wine, or hard liquor (for example, vodka, whiskey, or tequila) in your neighborhood, would he or she be caught by the police?

Definitely No Somewhat No Somewhat Yes Definitely Yes

If a kid carried a handgun in your neighborhood would he or she be caught by the police?

Definitely No Somewhat No Somewhat Yes Definitely Yes

COMMUNITY: Perceived Availability of Drugs

If you wanted to get some cigarettes, how easy would it Very hard be for you to get some?

Very hard Sort of Hard Sort of easy Very easy

If you wanted to get some beer, wine, or hard liquor (for example, vodka, whiskey, or tequila), how easy would it be for you to get some?

Very hard Sort of Hard Sort of easy Very easy

If you wanted to get a drug like cocaine, LSD, or amphetamines, how easy would it be for you to get some?

Very hard Sort of Hard Sort of easy Very easy

If you wanted to get some marijuana, how easy would it Very hard be for you to get some?

Sort of Hard Sort of easy Very easy

COMMUNITY: Perceived Availability of Handguns

If you wanted to get a handgun, how easy would it be for you to get one?

Very hard Sort of Hard Sort of easy Very easy

COMMUNITY: Rewards for Prosocial Involvement

My neighbors notice when I am doing a good job and let me know about it.

Definitely No Somewhat No Somewhat Yes Definitely Yes

There are people in my neighborhood who are proud of me when I do something well.

Definitely No Somewhat No Somewhat Yes Definitely Yes

There are people in my neighborhood who encourage me to do my best.

Definitely No Somewhat No Somewhat Yes Definitely Yes

FAMILY: Poor Family Management

My parents ask if I've gotten my homework done. Definitely No

Somewhat No Somewhat Yes Definitely Yes

Would your parents know if you did not come home on

time?

Definitely No Somewhat No Somewhat Yes

Somewhat Yes Definitely Yes

The rules in my family are clear Definitely No

Somewhat No Somewhat Yes Definitely Yes When I am not at home, one of my parents knows

where I am and who I am with.

Definitely No Somewhat No Somewhat Yes **Definitely Yes**

If you drank some beer or wine or liquor (for example,

vodka, whiskey, or gin) without your parents' permission, would you be caught by your parents? Definitely No Somewhat No Somewhat Yes **Definitely Yes**

My family has clear rules about alcohol and drug use.

Definitely No Somewhat No Somewhat Yes **Definitely Yes**

If you carried a handgun without your parents' permission, would you be caught by your parents? Definitely No Somewhat No Somewhat Yes **Definitely Yes**

If you skipped school would you be caught by your parents?

Definitely No Somewhat No Somewhat Yes **Definitely Yes**

FAMILY: Family Conflict

People in my family often insult or yell at each other.

Definitely No Somewhat No Somewhat Yes **Definitely Yes**

We argue about the same things in my family over and

over.

Definitely No Somewhat No Somewhat Yes **Definitely Yes**

People in my family have serious arguments. **Definitely No**

Somewhat No Somewhat Yes **Definitely Yes**

FAMILY: Family History of Antisocial Behavior

Has anyone in your family ever had severe alcohol or drug problems?

No Yes

Have any of your brothers or sisters ever:

drunk beer, wine, or hard liquor (for example, vodka,

whiskey, or tequila)?

No Yes

I don't have any brothers or

sisters

smoked marijuana?

Yes

I don't have any brothers or

sisters

smoked cigarettes?

Yes

I don't have any brothers or

sisters

taken a handgun to school?

Yes

I don't have any brothers or

sisters

been suspended or expelled from school? No

Yes

I don't have any brothers or

sisters

used a vape product (e-cigarettes, vape pens, mods, or

pod vapes like JUUL or Puff Bars)?

No

Yes

I don't have any brothers or

sisters

About how many adults (over 21, if any) have you known personally who in the past year have:

used marijuana, crack cocaine, or other drugs?

1 adult 2 adults

3- 4 adults

5 + adults

sold or dealt drugs?

2 adults 3- 4 adults 5 + adults

1 adult

0

done other things that could get them in trouble with the police, like stealing, selling stolen goods, mugging

or assaulting others, etc?

1 adult 2 adults 3- 4 adults 5 + adults

gotten drunk or high? 0

1 adult 2 adults 3- 4 adults 5 + adults

FAMILY: Parental Attitudes Favorable Toward Drug Use

How wrong do your parents feel it would be for YOU to:

drink beer, wine, or hard liquor (for example, vodka, whiskey, or tequila) regularly? Very wrong Wrong

A little bit wrong Not wrong at all

smoke cigarettes? Very wrong

Wrong

A little bit wrong Not wrong at all

smoke marijuana? Very wrong

Wrong

A little bit wrong Not wrong at all

FAMILY: Parental Attitudes Favorable Toward Antisocial Behavior

steal anything worth more than \$5? Very wrong

Wrong

A little bit wrong Not wrong at all

draw graffiti, or write things, or draw pictures on buildings or other property (without the owner's

permission)?

Very wrong Wrong

A little bit wrong Not wrong at all

pick a fight with someone? Very wrong

Wrong

A little bit wrong Not wrong at all

FAMILY: Attachment

Do you feel very close to your mother?

Definitely No

Somewhat No Somewhat Yes Definitely Yes

Do you share your thoughts and feeling with your

mother?

Definitely No Somewhat No Somewhat Yes Definitely Yes

Do you feel very close to your father?

Definitely No

Somewhat No Somewhat Yes Definitely Yes

Do you share your thoughts and feeling with your

father?

Definitely No Somewhat No Somewhat Yes Definitely Yes

FAMILY: Opportunities for Prosocial Involvement

My parents ask me what I think before most family

décisions affecting me are made.

Definitely No Somewhat No Somewhat Yes **Definitely Yes**

If I had a personal problem, I could ask my mom or dad

for help.

Definitely No Somewhat No Somewhat Yes **Definitely Yes**

Definitely No

My parents give me lots of chances to do fun things with them.

Somewhat No Somewhat Yes **Definitely Yes**

FAMILY: Rewards for Prosocial Involvement

Do you enjoy spending time with your mother? **Definitely No**

> Somewhat No Somewhat Yes **Definitely Yes**

Do you enjoy spending time with your father? **Definitely No**

Somewhat No Somewhat Yes **Definitely Yes**

My parents notice when I am doing a good job and let

me know about it.

Never or almost never

Sometimes

Often

All the time

How often do your parents tell you they're proud of you Never or almost never

for something you've done?

Sometimes

Often

All the time

SCHOOL: Academic Failure

Are your school grades better than the grades of most

students in your class?

Definitely No Somewhat No Somewhat Yes Definitely Yes

Putting them all together, what were your grades like

last year?

Mostly F's
Mostly D's
Mostly C's
Mostly B's
Mostly A's

SCHOOL: Little Commitment to School

How often do you feel that the school work you are

assigned is meaningful and important?

Almost Always

Often

Sometimes Seldom Never

How important do you think the things you are learning

in school are going to be for your later life?

Very important
Quite important
Fairly important
Slightly important
Not at all important

How interesting are most of your courses to you?

Very interesting & stimulating

Quite interesting Fairly interesting Slightly interesting Not at all interesting

Now, thinking back over the past year in school, how often did you...

enjoy being in school?

Never Seldom Sometimes Often

Almost Always

hate being in school? Never

Seldom Sometimes Often

Almost Always

try to do your best work in school?

Never

Seldom Sometimes Often

Almost Always

During the last four weeks, how many (if any) whole days of school have you missed because you skipped or

None 1

2

3 4-5

6-10

11 or more days

SCHOOL: Opportunities for Prosocial Involvement

In my school, students have lots of chances to help decide things like class activities and rules.

Definitely No Somewhat No Somewhat Yes Definitely Yes

Teachers ask me to work on special classroom projects.

Definitely No Somewhat No Somewhat Yes Definitely Yes

There are lots of chances for students in my school to get involved in sports, clubs, and other school activities outside of class.

Definitely No Somewhat No Somewhat Yes Definitely Yes

There are lots of chances for students in my school to talk with a teacher one-on-one.

Definitely No Somewhat No Somewhat Yes Definitely Yes I have lots of chances to be part of class discussions or activities.

Definitely No Somewhat No Somewhat Yes Definitely Yes

SCHOOL: Rewards for Prosocial Involvement

My teachers notice when I am doing a good job and lets Definitely No

me know about it.

Somewhat No

Somewhat Yes Definitely Yes

I feel safe at my school. Definitely No

Somewhat No Somewhat Yes Definitely Yes

The school lets my parents know when I have done

something well.

Definitely No Somewhat No Somewhat Yes Definitely Yes

My teachers praise me when I work hard in school. Definitely No

Somewhat No Somewhat Yes Definitely Yes

PEER-INDIVIDUAL: Rebelliousness

I do the opposite of what people tell me, just to get

them mad.

Very False

Somewhat False Somewhat True

Very True

I like to see how much I can get away with. Very False

Somewhat False Somewhat True

Very True

I ignore the rules that get in my way.

Very False Somewhat False Somewhat True Very True

PEER-INDIVIDUALS: Early Initiation of Drug Use

If ever, how old were you when you first:

used marijuana (grass, pot, cannabis, weed) or hashish (hash, hash oil)?	Never 10 or younger 11 12 13 14 15 16 17 or older
smoked a cigarette, even just a puff?	Never 10 or younger 11 12 13 14 15 16 17 or older
had more than a sip or two of beer, wine, or hard liquor (for example, vodka, whiskey, or tequila)?	Never 10 or younger 11 12 13 14 15 16 17 or older

began drinking alcoholic beverages regularly, that is, at least once or twice a month?

Never

10 or younger

11

12

13

14

15 16

17 or older

PEER-INDIVIDUALS: Early Initiation of Antisocial Behavior

If ever, how old were you when you first:

got suspended from school?

Never

10 or younger

11

12

13

14 15

16

17 or older

got arrested?

Never

10 or younger

11

12

13

14

15

16

17 or older

carried a handgun? Never

10 or younger

11

12

13 14

15 16

17 or older

attacked someone with the idea of seriously hurting

them?

Never

10 or younger

11

12

13

14 15

16

17 or older

PEER-INDIVIDUALS: Favorable Attitudes Toward Antisocial Behavior

How wrong do you think it is for someone your age to...

take a handgun to school? Very Wrong

Wrong

A Little Bit Wrong Not Wrong at All

steal anything worth more than \$5? Very Wrong

Wrong

A Little Bit Wrong Not Wrong at All

pick a fight with someone? Very Wrong

Wrong

A Little Bit Wrong Not Wrong at All attack someone with the idea of seriously hurting

them?

Very Wrong Wrong

A Little Bit Wrong Not Wrong at All

stay away from school all day when their parents think

they are at school?

Very Wrong Wrong

A Little Bit Wrong Not Wrong at All

PEER-INDIVIDUALS: Favorable Attitudes Toward Drug Use

How wrong do you think it is for someone your age to:

drink beer, wine, or hard liquor (for example, vodka,

whiskey, or tequila) regularly?

Very Wrong Wrong

A Little Bit Wrong Not Wrong at All

smoke cigarettes? Very Wrong

Wrong

A Little Bit Wrong Not Wrong at All

smoke marijuana? Very Wrong

Wrong

A Little Bit Wrong Not Wrong at All

use LSD, cocaine, amphetamines or another illegal

drug?

Very Wrong

Wrong

A Little Bit Wrong Not Wrong at All

PEER-INDIVIDUALS: Perceived Risks of Drug Use

How much do you think people risk harming themselves (physically or in other ways) if they:

Smoke one or more packs of cigarettes per day? No Risk Slight Risk Moderate Risk

Great Risk

No Risk Try marijuana once or twice?

Slight Risk Moderate Risk **Great Risk**

Smoke marijuana regularly? No Risk

> Slight Risk Moderate Risk **Great Risk**

No Risk

Take one or two drinks of an alcoholic beverage (beer, wine, liquor) nearly every day.

Slight Risk Moderate Risk **Great Risk**

PEER-INDIVIDUALS: Interaction with Antisocial Peers

Think of you four best friends (the friends you feel closest to). In the past year (12 months), how many of your best friends have:

been suspended from school? 0 1 2 3 4 carried a handgun? 0 2 3

sold illegal drugs?	0 1 2 3 4
stolen or tried to steal a motor vehicle such as a car or motorcycle?	0 1 2 3 4
been arrested?	0 1 2 3 4
dropped out of school?	0 1 2 3 4

PEER-INDIVIDUALS: Friends' Use of Drugs

Think of you four best friends (the friends you feel closest to). In the past year (12 months), how many of your best friends have:

smoked cigarettes?	0
	2 3 4
tried beer, wine, or hard liquor (for example, vodka, whiskey, or tequila) when their parents didn't know about it?	0 1 2 3 4

used marijuana?	0
	1
	2
	3
	4
used LSD, cocaine, amphetamines or another illegal	0
drugs?	1
	2
	3
	4

PEER-INDIVIDUALS: Rewards for Antisocial Involvement

What are the chances you would be seen as cool if you:

smoked cigarettes?	No or Very Little Chance Little Chance Some Chance Pretty Good Chance Very Good Chance
began drinking alcoholic beverages regularly, that is, at least once or twice a month?	No or Very Little Chance Little Chance Some Chance Pretty Good Chance Very Good Chance
smoked marijuana?	No or Very Little Chance Little Chance Some Chance Pretty Good Chance Very Good Chance
carried a handgun?	No or Very Little Chance Little Chance Some Chance Pretty Good Chance Very Good Chance

PEER-INDIVIDUALS: Gang Involvement scale

Have you ever belonged to a gang?

No, but would like to

No

Yes, in the past Yes, belong now

Yes, but would like to get out

PEER-INDIVIDUALS: Depressive Symptoms

Sometimes I think that life is not worth it.

Definitely No

Somewhat No Somewhat Yes Definitely Yes

At times I think I am no good at all.

Definitely No

Somewhat No Somewhat Yes Definitely Yes

All in all, I am inclined to think that I am a failure. Definitely No

Somewhat No Somewhat Yes Definitely Yes

In the past year have you felt depressed or sad MOST

days, even if you felt OK sometimes?

Definitely No Somewhat No Somewhat Yes Definitely Yes

PEER-INDIVIDUALS: Belief in Moral Order

I think sometimes it's okay to cheat at school. Definitely No

Somewhat No Somewhat Yes Definitely Yes

I think it is okay to take something without asking if you

can get away with it.

Definitely No Somewhat No Somewhat Yes Definitely Yes It is all right to beat up people if they start the fight. Definitely No

Somewhat No Somewhat Yes Definitely Yes

It is important to be honest with your parents, even if they become upset or you get punished.

Definitely No Somewhat No Somewhat Yes Definitely Yes

PEER-INDIVIDUALS: Prosocial Involvement

If ever, how many times in the past year (12 months) have you:

participated in clubs, organizations and activities at school?	Never 1 or 2 times 3-5 6-9 10-19 20-29 30-39 40+
done extra work on your own for school?	Never 1 or 2 times 3-5 6-9 10-19 20-29 30-39 40+
volunteered to do community service?	Never 1 or 2 times 3-5 6-9 10-19 20-29 30-39 40+

PEER-INDIVIDUALS: Rewards for Prosocial Involvement

What are the chances you would be seen as cool if you:

worked hard in school?

Very good chance
Pretty good chance
Some chance

Little chance

No or very little chance

defended someone who was being verbally abused at

school?

Very good chance Pretty good chance Some chance

Little chance No or very little chance

regularly volunteered to do community service? Very good chance

Pretty good chance

Some chance Little chance

No or very little chance

PEER-INDIVIDUALS: Interaction with Prosocial Peers

Think of your four best friends (the friends you feel closest to). In the past year (12 months), how many (if any) of your best friends have:

participated in clubs, organizations and activities at school?

1

2

3

4

made a commitment to stay drug-free?

1

2

3

4

tried to do well in school?	0 1
	2 3
	4
liked school?	0
	1
	2
	3
	4
regularly attended religious services?	0
	1
	2
	3
	4

DRUG USE

Think back over the last two weeks; if any, how many times have you had five or more alcoholic drinks in a row?

None Once Twice

3-5 times 6-9 times

10 or more times

If ever, how old were you when you first: used a vape product (e-cigarettes, vape pens, mods, or pod vapes like JUUL or Puff Bars)?

Never

10 or younger

11

12

13

14

15 16

17 or older

On how many occasions (if any) have you had alcoholic beverages (beer, wine or hard liquor) to drink in your lifetime - more than just a few sips?	0 occasions 1-2 3-5 6-9 10-19 20-39 40 or more
On how many occasions (if any) have you had beer, wine or hard liquor during the past 30 days?	0 occasions 1-2 3-5 6-9 10-19 20-39 40 or more
If ever, how old were you when you first: used marijuana (grass, pot, cannabis, weed) or hashish (hash, hash oil).	Never 10 or younger 11 12 13 14 15 16 17 or older
On how many occasions (if any) have you used marijuana (grass, pot) or hashish (hash, hash oil) during the past 30 days?	0 occasions 1-2 3-5 6-9 10-19 20-39 40+ above
If ever, how old were you when you first: used LSD (acid) or other hallucinogens (like PCP, mescaline, "shrooms" or psilocybin).	Never 10 or younger 11 12 13 14 15 16 17 or older

On how many occasions (if any) have you used LSD (acid) or other hallucinogens (like PCP, mescaline, "shrooms" or psilocybin during the past 30 days?	0 occasions 1-2 3-5 6-9 10-19 20-39 40+ above
If ever, how old were you when you first: used cocaine (like cocaine powder) or "crack" (cocaine in chunk or rock form)?	Never 10 or younger 11 12 13 14 15 16 17 or older
On how many occasions (if any) have you used cocaine (like cocaine powder) or "crack" (cocaine in chunk or rock form) during the past 30 days?	0 occasions 1-2 3-5 6-9 10-19 20-39 40+ above
If ever, how old were you when you first: sniffed glue, breathed the contents of an aerosol spray can, or inhaled other gases or sprays, in order to get high?	Never 10 or younger 11 12 13 14 15 16 17 or older
On how many occasions (if any) have you sniffed glue, breathed the contents of an aerosol spray can, or inhaled other gases or sprays, in order to get high during the past 30 days?	0 occasions 1-2 3-5 6-9 10-19 20-39 40+ above

If ever, how old were you when you first: used methamphetamines (meth, speed, crank, crystal meth)?	Never 10 or younger 11 12 13 14 15 16 17 or older
On how many occasions (if any) have you used methamphetamines (meth, speed, crank, crystal meth) in the past 30 days?	0 occasions 1-2 3-5 6-9 10-19 20-39 40+ above
If ever, how old were you when you first: used prescription stimulants or amphetamines (such as Adderall, Ritalin, or Dexedrine) without a doctor telling you to take them?	Never 10 or younger 11 12 13 14 15 16 17 or older
On how many occasions (if any) have you used prescription stimulants or amphetamines (such as Adderall, Ritalin, or Dexedrine) without a doctor telling you to take them, during the past 30 days?	0 occasions 1-2 3-5 6-9 10-19 20-39 40+ above

If ever, how old were you when you first: used prescription sedatives including barbiturates or sleeping pills (such as phenobarbital, Tuinal, Seconal, Ambien, Lunesta, or Sonata) without a doctor telling you to take them?

Never 10 or younger 11 12

17 or older

On how many occasions (if any) have you used prescription sedatives including barbiturates or sleeping pills (such as phenobarbital, Tuinal, Seconal, Ambien, Lunesta, or Sonata) without a doctor telling you to take them, during the past 30 days?

0 occasions

1-2 3-5

6-9

10-19 20-39 40+ above

If ever, how old were you when you first: used prescription tranquilizers (such as Librium, Valium, Xanax, Ativan, Soma, or Klonopin) without a doctor telling you to take them?

Never

10 or younger

11

12

13

14 15

16

17 or older

On how many occasions (if any) have you used prescription tranquilizers (such as Librium, Valium, Xanax, Ativan, Soma, or Klonopin) without a doctor telling you to take them, during the past 30 days?

0 occasions

1-2

3-5

6-9

10-19

20-39

40+ above

If ever, how old were you when you first: used narcotic prescription drugs (such as OxyContin, methadone, morphine, codeine, Demerol, Vicodin, Percocet) without a doctor telling you to take them?	Never 10 or younger 11 12 13 14 15 16 17 or older
On how many occasions (if any) have you used narcotic prescription drugs (such as OxyContin, methadone, morphine, codeine, Demerol, Vicodin, Percocet) without a doctor telling you to take them, during the past 30 days?	0 occasions 1-2 3-5 6-9 10-19 20-39 40+ above
If ever, how old were you when you first: used heroin?	Never 10 or younger 11 12 13 14 15 16 17 or older
On how many occasions (if any) have you used heroin during the past 30 days?	0 occasions 1-2 3-5 6-9 10-19 20-39 40+ above
On how many occasions (if any) have you used steroids or anabolic steroids (such as Anadrol, Oxandrin, Durabolin, Equipoise, or Depotesterone) in the past 30 days?	0 occasions 1-2 3-5 6-9 10-19 20-39 40+ above

During the past 30 days, how did you use marijuana? (Mark ALL that apply.)

I did not use marijuana during the past 30 days

I smoked it

l ate it (in an edible, candy, tincture or other food)

I used a vaporizer

I dabbed it

I used it in some other way

ANTISOCIAL BEHAVIOR

If ever, how many times in the past year (12 months) have you:

been suspended from school?	Never 1 or 2 times 3-5 6-9 10-19 20-29 30-39 40+
carried a handgun?	Never 1 or 2 times 3-5 6-9 10-19 20-29 30-39 40+
sold illegal drugs?	Never 1 or 2 times 3-5 6-9 10-19 20-29 30-39 40+

stolen or tried to steal a motor vehicle such as a car or motorcycle?	Never 1 or 2 times 3-5 6-9 10-19 20-29 30-39 40+
been arrested?	Never 1 or 2 times 3-5 6-9 10-19 20-29 30-39 40+
attacked someone with the idea of seriously hurting them?	Never 1 or 2 times 3-5 6-9 10-19 20-29 30-39 40+
been drunk or high at school	Never 1 or 2 times 3-5 6-9 10-19 20-29 30-39 40+

taken a handgun to school

1 or 2 times
3-5
6-9
10-19
20-29

30-39 40+

NEED FOR TREATMENT

In the past 12 months have you spent more time using No alcohol than you intended? Yes Don't Use In the past 12 months have you spent more time using No drugs than you intended? Yes Don't Use In the past 12 months, have you neglected some of No your usual responsibilities because of using alcohol? Yes Don't Use In the past 12 months, have you neglected some of No your usual responsibilities because of using drugs? Yes Don't Use In the past 12 months, have you wanted to cut down on No your alcohol use? Yes Don't Use In the past 12 months, have you wanted to cut down on No your drug use? Yes Don't Use In the past 12 months, has anyone objected to your No alcohol use? Yes Don't Use In the past 12 months, has anyone objected to your No drug use? Yes Don't Use In the past 12 months, did you frequently find yourself

thinking about using alcohol?

No Yes

Don't Use

In the past 12 months, did you frequently find yourself thinking about using drugs?

No Yes

Don't Use

In the past 12 months, did you use alcohol to relieve feelings such as sadness, anger, or boredom?

No

Yes

Don't Use

In the past 12 months, did you use drugs to relieve feelings such as sadness, anger, or boredom?

No Yes

Don't Use

UTAH STATE BOARD OF EDUCATION: SCHOOL CLIMATE **ITEMS**

How worried, if at all, are you about the possibility of each of the following things happening at your school?

Getting bullied Not at all worried

> Not too worried Somewhat worried

Very worried

Gun violence or active shooter situation Not at all worried

> Not too worried Somewhat worried

Very worried

Suicide by a student Not at all worried

> Not too worried Somewhat worried

Very worried

Gang activity Not at all worried

> Not too worried Somewhat worried

Very worried

Appendix F: Item Dictionary

Students using alcohol or drugs Not at all worried

Not too worried Somewhat worried

Very worried

Earthquake/fire Not at all worried

Not too worried Somewhat worried

Very worried

How safe do you feel in each of the following areas at your school (before and after school)?

Playgrounds or fields Very unsafe

Somewhat unsafe Somewhat safe

Very safe

Lunchroom/Cafeteria Very unsafe

Somewhat unsafe Somewhat safe

Very safe

Classrooms Very unsafe

Somewhat unsafe Somewhat safe

Very safe

Bathrooms Very unsafe

Somewhat unsafe Somewhat safe

Very safe

Parking lots Very unsafe

Somewhat unsafe Somewhat safe

Very safe

Stairs and hallways Very unsafe

Somewhat unsafe Somewhat safe

Very safe

On the school bus

Very unsafe Somewhat unsafe Somewhat safe Very safe

PARENTS EMPOWERED QUESTIONS

My parents have set clear rules and expectations with me about NOT drinking ANY alcohol. Definitely No Somewhat No Somewhat Yes Definitely Yes

During the past year (12 months), how often have you talked with at least one of your parents about the rules and expectations of NO alcohol use?

At least once a month Every 2 to 3 months Every 4 to 6 months

A few times in the past year Talked, but not in the past year

Never Never

DRUG FREE COMMUNITIES QUESTIONS

During the past 12 months, have you talked with at least one of your parents about the dangers of tobacco, alcohol, or drug abuse? (Choose all that apply).

Yes, tobacco use Yes, alcohol use Yes, drug use

How much do you think people risk harming themselves (physically or in other ways) if they: have five or more drinks once or twice each weekend?

No Risk Slight Risk Moderate Risk Great Risk

How much do you think people risk harming themselves (physically or in other ways) if they: Have five or more drinks of an alcoholic beverage once or twice a week? No Risk Slight Risk Moderate Risk Great Risk

How much do you think people risk harming themselves (physically or in other ways) if they: Smoke marijuana once or twice a week?

No Risk Slight Risk Moderate Risk Great Risk How much do you think people risk harming themselves (physically or in other ways) if they: Use prescription drugs that are not prescribed for them?

No Risk Slight Risk Moderate Risk Great Risk

How much do you think people risk harming themselves (physically or in other ways) if they: Smoke 1-5 cigarettes per day?

No Risk Slight Risk Moderate Risk Great Risk

How wrong do your parents feel it would be for YOU to: have one or two drinks of an alcoholic beverage nearly every day?

Very wrong Wrong A little bit wrong Not wrong at all

How wrong do your parents feel it would be for YOU to: Very wrong use prescription drugs not prescribed to you?

Wrong

Wrong
A little bit wrong
Not wrong at all

How wrong do your friends feel it would be for you to: have one or two drinks of an alcoholic beverage nearly every day? Very wrong Wrong

A little bit wrong Not wrong at all

How wrong do your friends feel it would be for you to: smoke tobacco?

Very wrong Wrong

A little bit wrong Not wrong at all

How wrong do your friends feel it would be for you to: smoke marijuana?

Very wrong Wrong

A little bit wrong Not wrong at all

How wrong do your friends feel it would be for you to: use prescription drugs not prescribed to you?

Very wrong Wrong

A little bit wrong Not wrong at all During the past 30 days, did you drive a car or other vehicle when you had been drinking alcohol? If so, how many times?

I do not drive 0 times 1 time 2 or 3 times

4 or 5 times 6 or more times

SCHOOL SAFETY

During the past 30 days, on how many days (if any) did you not go to school because you felt you would be unsafe at school or on the way to or from school?

0 days 1 day 2-3 days 4-5 days 6 or more days

During the past 12 months, how often (if at all) have you been picked on or bullied by a student ON SCHOOL PROPERTY?

0 days 1 day 2-3 days 4-5 days

6 or more days

During the past 12 months, how often (if at all), have you been threatened or harassed over the internet, by e-mail, or by someone using a cell phone?

0 times 1 time 2 or 3 times 4 or 5 times

6 or more times

PLACES OF ALCOHOL

During the past year, did you drink alcohol at any of the At my home or someone else's following places? (Mark the number of times for each.)

home without any parent permission

At my home with my parents' permission

At someone else's home with their parents' permission

In another place

In a car

At or near school

Someplace outside of town (for example, on public lands, in the desert, in a campground, etc.)

SOCIAL DETERMINANTS OF HEALTH

This past year, did you experience any of the following? (Mark ALL that apply.)

One or more people living in my home lost their job

I had to move or change homes in the past year

Skipped one or more meals because my family didn't have enough money to buy food

I had difficulty keeping up with schoolwork because I didn't have access to a reliable computer or internet service

I did not have a quiet place at home to study

None of these

HEALTH DEPARTMENT QUESTIONS

How often do you wear a seatbelt when riding in a car driven by someone else?

Never Rarely Sometimes Most of the time **Always**

My parents expect me to eat dinner at home with my family.	Definitely No Somewhat No Somewhat Yes Definitely Yes
During the past 12 months, about how many days of school did you miss because of your asthma?	I do not have asthma 0 days 1 to 3 days 4 to 9 days 10 to 12 days 13 or more days
During the past 7 days, on how many days were you physically active for a total of at least 60 minutes per day? (Add up all the time you spent in any kind of physical activity that increased your heart rate and made you breathe hard some of the time.)	0 days 1 day 2 days 3 days 4 days 5 days 6 days 7 days
On an average school day, how many hours do you use an electronic device for something that is not school work? (Count time spent on things such as Xbox, PlayStation, texting, YouTube, Instagram, Facebook, or other social media.):	Less than 1 hour per day 1 hour per day 2 hours per day 3 hours per day 4 hours per day 5 or more hours per day
In a typical week, how many days do you walk, ride your bike or scooter (non-motorized), or skateboard to and from school?	No days 1 2 3 4 5
Has a doctor or nurse ever told you that you have asthma?	No Yes
Do you still have asthma	No Yes
During the past 12 months, did you have an episode of asthma or an asthma attack?	No Yes

An asthma action plan, or asthma management plan, is a form with instructions about when to change the amount or type of medicine, when to call the doctor for advice, and when to go to the emergency room. Has a doctor or other health professional EVER given you a written asthma action plan?

I do not have asthma

Yes No

Not sure

Do you have diabetes?

No Yes

Has a doctor or other health professional EVER given you a written diabetes care plan to help manage your diabetes in school?

I do not have diabetes

Yes No

Not sure

During the past 12 months, about how many days of school did you miss because of your diabetes?

I do not have diabetes

0 days 1 to 3 days 4 to 9 days 10 to 12 days 13 or more days

During the past 12 months, did you ever feel so sad or hopeless almost every day for two weeks or more in a row that you stopped doing some usual activities?

No

Yes

During the past 12 months, did you ever seriously consider attempting suicide?

No Yes

During the past 12 months, did you make a plan about how you would attempt suicide?

No

Yes

During the past 12 months, how many times (if any)

did you actually attempt suicide?

0 times 1 time

2 to 3 times

4 to 5 times

6 or more times

In the past seven days, I have felt left out.

Never Rarely

Sometimes

Often Always In the past seven days, I have felt that people barely know me.

Never Rarely

Sometimes

Often **Always**

In the past seven days, I have felt isolated from others.

Never Rarely Sometimes

Often **Always**

In the past seven days, I have felt that people are

around me but not with me.

Never Rarely Sometimes Often

Always

Grid

How tall are you without your shoes on? Grid

How much do you weigh without your shoes on?

During the past 30 days, did you drive a car or other vehicle when you were talking on a cell phone? If so, on how many days?

0 times 1 time

2 or 3 times 4 or 5 times 6 or more times

During the past 30 days, did you text or e-mail while driving a car or other vehicle? If so, on how many days?

0 times 1 time

2 or 3 times 4 or 5 times 6 or more times

How are guns and bullets stored in your home?

We don't have any guns or

bullets.

Unlocked and in plain sight Locked or hidden, but I know

how to access them.

Locked or hidden, and I DON'T know how to access them

Don't know

SELF-HARM QUESTIONS

In the past 12 months, have you ever done something to purposefully hurt yourself without wanting to die, such as cutting or burning yourself on purpose? If so, how many times did you do so?

0 times 1 time

2 or 3 times

4 or 5 times

6 or more times

TOBACCO QUESTIONS

Have you ever tried:

cigarettes, even just one puff?	No Yes
cigars, cigarillos, or little cigars, even a puff?	No Yes
tobacco in a hookah or waterpipe?	No Yes
vape products containing nicotine (such as e-cigarettes, vape pens, mods, or pod vapes like JUUL or Puff Bars)?	No Yes
vape products containing marijuana?	No Yes
chewing tobacco, snuff, dip, or snus (moist smokeless tobacco usually sold in small pouches)?	No Yes
nicotine pouches like Zyn, On, and Velo?	No Yes
nicotine lozenges (small hard or soft tablest containing nicotine that slowly dissolve in the mouth such as Velo or Rouge)?	No Yes

During the past 30 days, on how many days did you:

Smoke tobacco in a hookah or waterpipe? O days 1 or 2 days 3 to 5 days 6 to 9 days 10 to 19 days 20 to 29 days all 30 days Use vape products containing nicotine (such as e-cigarettes, vape pens, mods, or pod vapes like JULL or Puff Bars)? O days 1 or 2 days 3 to 5 days 6 to 9 days 1 or 2 days 3 to 5 days 6 to 9 days 10 to 19 days 20 to 29 days all 30 days Use vape products containing marijuana? O days 1 or 2 days 3 to 5 days 6 to 9 days 1 or 2 days 3 to 5 days 6 to 9 days 1 or 2 days 3 to 5 days 6 to 9 days 1 or 2 days 3 to 5 days 6 to 9 days 1 or 2 days 3 to 5 days 6 to 9 days 1 or 2 days 3 to 5 days 6 to 9 days 1 or 2 days 3 to 5 days 6 to 9 days 1 or 10 to 19 days 10 to 19 days	Smoke cigarettes?	0 days 1 or 2 days 3 to 5 days 6 to 9 days 10 to 19 days 20 to 29 days all 30 days
1 or 2 days 3 to 5 days 6 to 9 days 10 to 19 days 20 to 29 days all 30 days use vape products containing nicotine (such as e-cigarettes, vape pens, mods, or pod vapes like JULL or Puff Bars)? 0 days 1 or 2 days 3 to 5 days 6 to 9 days 10 to 19 days 20 to 29 days all 30 days use vape products containing marijuana? 0 days 1 or 2 days 3 to 5 days 6 to 9 days 1 or 2 days 3 to 5 days 6 to 9 days 1 or 2 days 3 to 5 days 1 or 2 days 3 to 5 days 6 to 9 days 1 or 2 days 3 to 5 days 6 to 9 days 10 to 19 days 20 to 29 days	Smoke cigars, cigarillos, or little cigars?	1 or 2 days 3 to 5 days 6 to 9 days 10 to 19 days 20 to 29 days
e-cigarettes, vape pens, mods, or pod vapes like JULL or 1 or 2 days 3 to 5 days 6 to 9 days 10 to 19 days 20 to 29 days all 30 days use vape products containing marijuana? 0 days 1 or 2 days 3 to 5 days 6 to 9 days 3 to 5 days 6 to 9 days 10 to 19 days 20 to 29 days	Smoke tobacco in a hookah or waterpipe?	1 or 2 days 3 to 5 days 6 to 9 days 10 to 19 days 20 to 29 days
1 or 2 days 3 to 5 days 6 to 9 days 10 to 19 days 20 to 29 days	e-cigarettes, vape pens, mods, or pod vapes like JULL or	1 or 2 days 3 to 5 days 6 to 9 days 10 to 19 days 20 to 29 days
	use vape products containing marijuana?	1 or 2 days 3 to 5 days 6 to 9 days 10 to 19 days 20 to 29 days

Use chewing tobacco, snuff, dip, or snus (moist smokeless tobacco usually sold in small pouches)?

0 days 1 or 2 days 3 to 5 days

6 to 9 days 10 to 19 days 20 to 29 days

all 30 days

Use nicotine pouches like Zyn, On, and Velo?

0 days 1 or 2 days 3 to 5 days 6 to 9 days 10 to 19 days 20 to 29 days all 30 days

use nicotine lozenges (small hard or soft tablets containing nicotine that slowly dissolve in the mouth such as Velo or Rogue)?

1 or 2 days 3 to 5 days 6 to 9 days 10 to 19 days 20 to 29 days all 30 days

0 days

How frequently (if ever) have you smoked cigarettes during the past 30 days?

Not at all

Less than one cigarette per day One to five cigarettes per day About one-half pack per day About one pack per day

About one and one-half packs

per day

Two packs or more per day

How much do you think people risk harming themselves (physically or in other ways) if they: use vape products such as e-cigarettes, vape pens, mods, or pod vapes like JUUL or Puff Bars? No risk Slight risk Moderate risk Great risk How old were you when you first used a vape product (e-cigarettes, vape pens, mods, or pod vapes like JUUL or Puff Bars)?

Never

10 or younger

11

12

13

14 15

16

17 or older

How wrong do your parents feel it would be for YOU to use vape products such as e-cigarettes, vape pens, mods, or pod vapes like JUUL or Puff Bars?

Very Wrong Wrong A little bit wrong Not wrong at all

If you smoked cigarettes or used vape products in the past 30 days, how did you usually get your own cigarettes or vape products? (CHOOSE ONLY ONE ANSWER FOR EACH TOBACCO TYPE- Vape Products, Regular Cigarettes)

- I did not use cigarettes or vape products (e-cigarettes, vape pens, mods, or pod vapes like JUUL) in the past 30 days
- I bought them in a store such as a convenience store, supermarket, discount store, or gas station
- I bought them at a tobacco smoke or vape shop
- I bought them at a vape shop
- I bought them on the internet or social media (such as Facebook, Instagram, or SnapChat)
- I gave someone else money to buy them for me
- I borrowed (or bummed) them from someone else
- A person 18 years old or older gave them to me
- I took them from a store or family member

I got them some other way

Do you think that you will try a cigarette soon?

I have already tried smoking cigarettes

No Yes If you have ever tried a tobacco product, which one did you try first?

I have never tried any tobacco product

Cigarettes

Cigars, cigarillos, or little cigars

Tobacco in a hookah or

waterpipe

Vape products such as

e-cigarettes, vape pens, or pod vapes like JUUL or Puff Bars? Chewing tobacco, snuff, or dip Nicotine pouches like Zyn, On, or

Velo Other

If you ever used vape products, such as e-cigarettes, vape pens, mods, or pod vapes like JUUL or Puff Bars, what flavor have you used most often?

I have never used vape product

Tobacco flavor Mint flavor Menthol flavor

Sweet, alcohol, or other flavor

If you smoked during the past 12 months, did you ever stop smoking for one day or longer because you were trying to quit smoking? I have not smoked in the past 12

months

Yes No

Do you think you will smoke a cigarette at any time during the next year?

Definitely yes Probably yes Probably not Definitely not

If one of your best friends offered you a cigarette, would you smoke it?

Definitely yes Probably yes Probably not Definitely not

Do you think that people can get addicted to vape products (e-cigarettes, vape pens, mods, or pod vapes like JUUL or Puff Bars)?

Definitely yes Probably yes Probably not Definitely not

Do you usually vape with nicotine, without nicotine, or both?

I do not vape now

Not at all A little Somewhat

A lot

How much do you want to stop vaping? I do not vape now Not at all A little Somewhat A lot Do you think you will use a vape product, such as Definitely yes e-cigarettes, vape pens, mods, or pod vapes like JUUL Probably yes or Puff Bars at any time during the next year? Probably not Definitely not If one of your best friends offered you a vape product Definitely yes such as an e-cigarette, vape pen, or mod, would you Probably yes use it? Probably not Definitely not Do you think people can get addicted to using tobacco Definitely yes just like they can get addicted to using cocaine or Probably yes heroin? Probably not Definitely not Do you think young people who smoke cigarettes have Definitely yes more friends? Probably yes Probably not Definitely not Do you think that smoke from other people's cigarettes Definitely yes is harmful to you? Probably yes Probably not Definitely not During this school year, were you taught in any of your No classes about the dangers of tobacco use? Yes Not sure During the past 7 days, on how many days were you in 0 days the same room with someone who was smoking 1 or 2 days cigarettes? 3 or 4 days 5 or 6 days 7 days

During the past 7 days, one how many days did you ride in a car with someone who was smoking

cigarettes?

0 days 1 or 2 days 3 or 4 days 5 or 6 days 7 days

Does anyone who lives with you now: (Mark ALL that

apply)

Smoke cigarettes

Use vape products (e-cigarettes, vape pens, mods, or pod vapes like JUUL or Puff Bars)

like JUUL or Puff Bars)
Use other tobacco products
No one lives with me now who
uses any form of tobacco

In the past 30 days, how often have you seen or heard any advertising or campaigns against smoking?

Never Rarely Sometimes Often Very Often

MENTAL HEALTH QUESTIONS

During the past 30 days, about how often did you:

feel nervous? All of the time

Most of the time Some of the time A little of the time None of the time

feel hopeless?

All of the time

Most of the time Some of the time A little of the time None of the time

feel restless or fidgety?

All of the time

Most of the time Some of the time A little of the time None of the time feel so depressed that nothing could cheer you up? All of the time

> Most of the time Some of the time A little of the time None of the time

feel that everything was an effort? All of the time

> Most of the time Some of the time A little of the time None of the time

feel worthless? All of the time

> Most of the time Some of the time A little of the time None of the time

How often in the last thirty days (if at all) did you talk to an adult (parent, doctor, counselor, teacher, etc.) about feeling very sad, hopeless, or suicidal?

I have not felt this way in the

past 30 days

0 times 1 time 2 to 4 times 5 or more times

If you have felt very sad, hopeless, or suicidal in the past 30 days who did you talk to about it? (Mark all that apply)

I have not felt this way in the

past 30 days

I felt this way but did not talk to

anyone about it

Parent Friend//Peer Teacher Doctor School Nurse School Counselor

Therapist

Clergy (e.g. Bishop, Priest or Nun, Minister, Pastor)

Other Adult

Do you think it's okay to seek help and talk to a professional counselor, therapist, or doctor if you've been feeling very sad, hopeless, or suicidal?

Yes No

I think it's okay for other people

to seek help

but not for me to seek help

During the past 12 months, how many times (if any) did someone you were dating or going out with physically hurt you on purpose? (Count such things as being hit, slammed into something, or injured with an object or weapon.)

I did not date or go out with anyone during the past 12 months

0 times 1 time

2 or 3 times 4 or 5 times 6 or more times

ADDITIONAL QUESTIONS

I feel safe in my neighborhood. Definitely No

Somewhat No Somewhat Yes Definitely Yes

If you have a religious preference, choose one which you identify the most.

Protestant (such as Baptists, Presbyterians, or Lutherans)

Catholic

Another religion

Iewish

LDS (Mormon) No preference

Now think about all the students in your grade at your school. How many of them do you think...

smoke one or more cigarettes a day? None (0%)

Few (1-10%) Some (11-30%) Half or less (31-50%)

Half or less (31-50%) Half or more (51-70%)

Most (71-90%)

Almost All (91-100%)

drank alcohol sometime in the past month? None (0%)

Few (1-10%) Some (11-30%) Half or less (31-50%) Half or more (51-70%)

Most (71-90%)

Almost All (91-100%)

used marijuana sometime in the past month? None (0%)

Few (1-10%) Some (11-30%) Half or less (31-50%) Half or more (51-70%)

Most (71-90%)

Almost All (91-100%)

use an illegal drug in the past month (not including marijuana)?

None (0%) Few (1-10%) Some (11-30%) Half or less (31-50%) Half or more (51-70%)

Most (71-90%)

Almost All (91-100%)

used a vape product (e-cigarettes, vape pens, mods, or

pod vapes like JUUL or Puff Bars)?

None (0%) Few (1-10%) Some (11-30%) Half or less (31-50%) Half or more (51-70%)

Most (71-90%)

Almost All (91-100%)

If you wanted to get vape products such as e-cigarettes, vape pens, mods, or pod vapes like JUUL or Puff Bars, how easy would it be for you to get some?

Very hard Sort of Hard Sort of easy Very easy

During a typical week, how many days do all or most of 0 your family eat at least one meal together? 1

1

3

4

5 6

7

HONESTY QUESTIONS

How old were you when you first: used phenoxydine (pox, px, breeze)

Never

10 or younger

11

12

13 14

15

16

17 or older

On how many occasions (if any) have you used phenoxydine (pox, px, breeze) in the past 30 days?

0 occasions

1-2

3-5

6-9 10-19

20-39

40+ above

How honest were you in filling out this survey?

I was very honest

I was honest most of the time I was honest some of the time I was honest once in a while

I was not honest at all